

# **Appendix E-1**

## **Arica Solar Project Biological Resources Technical Report**

**BIOLOGICAL RESOURCES TECHNICAL REPORT  
ARICA SOLAR PROJECT  
RIVERSIDE COUNTY, CALIFORNIA**



Prepared for:  
Aspen Environmental Group  
5020 Chesebro Rd. Suite 200  
Agoura Hills, CA 91301

and

Clearway Energy  
100 California Street,  
Suite 400  
San Francisco, CA 94111

Prepared by:



370 Alabama Street, Suite A  
Redlands, CA 92373

*Submitted: May 2021*

# Table of Contents

<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Background	1
1.2	Purpose	1
1.3	Site Location	1
1.4	Project Summary	2
<b>2</b>	<b>SITE CHARACTERISTICS</b>	<b>2</b>
2.1	Regional Setting	2
2.2	Hydrology	3
2.3	Soils	3
2.4	Sand Transport System	3
2.5	Rainfall	4
2.6	Vegetation Communities	5
2.6.1	Sonoran Creosote Bush Scrub	5
2.6.2	Desert Saltbush Scrub	6
3.1	Special Status Species Definition	6
3.2	Wildlife Surveys	7
3.2.1	Desert Tortoise	7
3.2.2	Mohave Fringe-toed Lizard	8
3.2.3	Couch's Spadefoot Toad	8
3.2.4	Avian Species	9
3.2.5	Special Status Bat Species	10
3.2.6	Other Special Status Wildlife Species	10
3.2.7	Wildlife Cameras	10
3.3	Special Status Plants	10
<b>4</b>	<b>RESULTS</b>	<b>12</b>
4.1	Special Status Wildlife	12
4.1.1	Desert Tortoise: <i>ST, FT</i>	13
4.1.2	Mojave fringe-toed lizard: <i>SSC, BLMS</i>	14
4.1.3	Couch's Spadefoot Toad: <i>SSC, BLMS</i>	14

4.1.4	Western Burrowing Owl: SSC, BCC, BLMS.....	14
4.1.5	Golden Eagle: CFP, WL, BCC, BLMS.....	15
4.1.6	Le Conte's Thrasher: SSC .....	17
4.1.7	Prairie Falcon: WL, BCC.....	17
4.1.8	Loggerhead Shrike: SSC (nesting), BCC.....	18
4.1.9	Gila Woodpecker: CE, BLMS, BCC.....	18
4.1.10	California Horned Lark: WL .....	18
4.1.11	Black-tailed Gnatcatcher: WL .....	19
4.1.12	Sonora Yellow Warbler: SSC, BCC .....	19
4.1.13	Short Eared Owl: SSC.....	19
4.1.14	Ferruginous Hawk: WL, BCC .....	19
4.1.15	Swainson's Hawk: ST, BBC.....	20
4.1.16	American Peregrine Falcon: FP, BCC.....	20
4.1.17	Vaux's Swift: SSC.....	20
4.1.18	Mountain Plover: SSC, BCC.....	20
4.1.19	Northern Harrier: SSC.....	21
4.1.20	Yellow-breasted Chat: SSC .....	21
4.1.21	Crissal Thrasher: SSC.....	21
4.1.22	Elf Owl: BLMS, BCC.....	21
4.1.23	Other listed Avian Species .....	22
4.1.23	American Badger: SSC .....	23
4.1.24	Desert Kit Fox: CPF .....	24
4.1.25	Desert Bighorn Sheep BLMS.....	24
4.1.26	Burro Deer: CPGS.....	24
4.2	Wildlife Movement.....	27
4.3	Special Status Plant Species .....	29
4.3.1	Crucifixion Thorn: CRPR 2B.2.....	30
4.3.2	Glandular Ditaxis: CRPR 2B.2.....	30
4.3.3	California Ditaxis: CRPR 3.2 .....	30
4.3.4	Utah Milkvine: CRPR 4.2 .....	30
4.3.5	Desert Unicorn Plant: CRPR 4.3.....	31
4.3.6	Chaparral Sand Verbena: BLMS, CRPR 1B.1 .....	31



4.3.7	Harwood's Milkvetch: CRPR 2B.2.....	31
4.3.8	Abram's Spurge: CRPR 2B.2.....	31
4.3.9	Ribbed Cryptantha: CRPR 4.3 .....	32
4.3.10	Harwood's Eriastrum: CRPR 1B.2, BLMS .....	32
4.3.11	Jackass Clover: CRPR 2B.2 .....	32
4.3.12	Palmer's Jackass Clover: CRPR 2B.2 .....	32
4.3.13	Creosote Bush Rings: BLMS .....	32
4.3.14	Cacti, Yucca, and Native Trees .....	32
4.4	Invasive Weeds.....	33
5.0	REFERENCES.....	36

Table 1.	Adjacent and Nearby Land Uses .....	2
Table 2.	Rainfall Summary .....	4
Table 3.	Plant Reference Population Visits.....	11
Table 4.	Special-status Wildlife and Plant Survey Personnel and Dates .....	12
Table 5.	Regional Golden Eagle Surveys .....	16
Table 6.	Summary of Wildlife Camera Observations.....	29

Photo 1.	Vegetation community – Sonoran creosote bush scrub.....	45
Photo 2.	Vegetation community – Saltbush scrub .....	45
Photo 3.	Vegetation community – Desert dry wash woodland .....	46
Photo 4.	Active aeolian sand .....	46
Photo 5.	Mohave fringe-toed lizard.....	47
Photo 6.	Canid burrow with burrowing owl white wash.....	47
Photo 7.	Camera station – burro deer.....	48
Photo 8.	Camera station – bob cat.....	48
Photo 9.	Camera station – coyote .....	49
Photo 10.	Camera station - jackrabbit.....	49
Photo 11.	Crucifixion thorn.....	50
Photo 12.	Eriastrum harwoodii.....	50
Photo 13.	Astragalus harwoodii .....	51
Photo 14.	Cryptantha costata.....	51

Figure 1.	General Vicinity.....	53
Figure 2.	Hydrologic Units .....	54
Figure 3.	Soils.....	55
Figure 4.	Aeolian Sand Transport .....	56

<b>Figure 5. Vegetation Communities .....</b>	<b>57</b>
<b>Figure 6. Study Area.....</b>	<b>58</b>
<b>Figure 7. Noteworthy Reptile and Amphibian Observations .....</b>	<b>59</b>
<b>Figure 8. Desert Tortoise Conservation Areas and Linkages .....</b>	<b>60</b>
<b>Figure 9. Noteworthy Avian Observations .....</b>	<b>61</b>
<b>Figure 10. Golden Eagle Survey Results 2010-2015 .....</b>	<b>62</b>
<b>Figure 11. Noteworthy Mammal Observations.....</b>	<b>63</b>
<b>Figure 12. Wildlife Connectivity.....</b>	<b>64</b>
<b>Figure 13. Noteworthy Plant Observations .....</b>	<b>65</b>

**Appendix A – Summary of Survey Results 2019-2020**

**A-1 Daily Survey Summary**

**A-2 Noteworthy Reptile and Amphibian Observations**

**A-3 Noteworthy Avian Observations**

**A-4 Avian Counts Summary**

**A-5 Noteworthy Mammal Observations**

**A-6 Noteworthy Plant Observations**

**A-7 Invasive Plant Species Observations**

**Appendix B – Potential for Special Status Wildlife Species to Occur**

**Appendix C – Potential for Special Status Plant Species to Occur**

**Appendix D - Wildlife Species Observed Fall 2019-Spring 2020**

**Appendix E – Plant Species Observed Fall 2019-Spring 2020**

## List of Acronyms

amsl	above mean sea level
ACEC	Area of Critical Environmental Concern
BRTR	Biological Resources Technical Report
BBCS	Bird and Bat Conservation Strategy
BLM	Bureau of Land Management
CA-177	California Highway 177
Cal-IPC	California Invasive Plant Council
CDFW	California Department of Fish and Wildlife
CDFA	California Department of Food and Agriculture
CESA	California Endangered Species Act
CEC	California Energy Commission
CEQA	California Environmental Quality Act
CDFW	California Department of Fish and Wildlife
CNPS	California Native Plant Society
CNDDDB	California Natural Diversity Database
CRPR	California Rare Plant Rank
DFA	Development Focus Area
DRECP	Desert Renewable Energy Conservation Plan
FEIS	Final Environmental Impact Statement
FESA	Federal Endangered Species Act
GIS	Geographic Information Systems
GPS	Global Positioning System
I-10	Interstate 10
LUPA	Land Use Plan Amendment
NEPA	National Environmental Protection Act
NPS	National Park Service
NECO Plan	Northern and Eastern Colorado Desert Coordinated Management Plan
O&M	Operations and Maintenance
PV	Photovoltaic
ROW	Right of Way
SEZ	Solar Energy Zone
TCAs	Tortoise Conservation Areas
USFWS	US Fish and Wildlife Service

# **1 INTRODUCTION**

## **1.1 Background**

Arica Solar, LLC is proposing to develop Arica Solar Project (Project) near the Desert Center community of unincorporated Riverside County, California. The proposed Project site is located on Bureau of Land Management (BLM) managed lands. The Arica Solar Project is expected to generate up to 265 megawatts (MW) of renewable energy using photovoltaic (PV) panels and will connect to the existing Red Bluff substation.

## **1.2 Purpose**

This Biological Resources Technical Report (BRTR) provides a description of methods and results of biological resource surveys and investigations conducted in fall of 2019 and spring of 2020 for the Arica Solar Project as approved by BLM in the project memo for the Biological Resources Survey Work Plan (Aspen 2019).

The primary purpose of this report is to provide biological information that will be used as the foundation for impact assessments pursuant to the National Environmental Policy Act (NEPA) and California Environmental Quality Act (CEQA). The discussion included herein may also be used to support consultation between Bureau of Land Management (BLM) and U.S. Fish and Wildlife Service (USFWS) under the Federal Endangered Species Act (FESA), and any necessary incidental take authorization from the California Department of Fish and Wildlife (CDFW) with respect to the California Endangered Species Act (CESA).

## **1.3 Site Location**

The Project site is in unincorporated eastern Riverside County, California. It consists of approximately 2,000 acres of BLM-managed land. The Project site is situated within Chuckwalla Valley near the community of Desert Center, nearly halfway between the cities of Indio and Blythe, north of the Interstate-10 freeway, on the Sidewinder Well and Corn Spring 7.5-Minute U.S. Geological Survey topographic quadrangles.

The Project site is located within in the California Desert Conservation Area (CDCA) planning area, and within the southern Desert Tortoise Recovery Unit of the Northern and Eastern Colorado Desert Coordinated Management (NECO) Plan. The Project site is not located within any ACECs (Areas of Critical Environmental Concern), but Alligator Rock ACEC is almost 3 miles southwest, the Desert Lily Preserve ACEC is 3 miles north, and Joshua Tree National Park is 5 miles north of the Project site. Desert tortoise critical habitat is located south of the I-10 freeway, less than 2 miles south of the site and a multi-species linkage area is just west of the site.

The entirety of the Project site is located within the boundaries of the Riverside East Solar Energy Zone (SEZ) identified in the Solar Programmatic Environmental Impact Statement (EIS) approved by a Record of Decision signed by BLM on October 12, 2012. Additionally, the Project site is within the Chuckwalla Valley ecoregion subarea of the Desert Renewable Energy Conservation Plan (DRECP) area. The DRECP identifies the federal lands in and around the Project site in the Land Use Plan Amendment (LUPA) and

Final Environmental Impact Statement (FEIS) as a Development Focus Area (DFA), as approved by a Record of Decision signed by BLM on September 14, 2016.

## 1.4 Project Summary

Arica Solar, LLC, a subsidiary of Clearway Energy Group LLC, has proposed to construct and operate the Arica Solar Project on land administered by BLM, on approximately 2,000 acres. The Project solar project site would be adjacent to one another concurrently proposed project (Victory Pass), also proposed by Clearway Energy that will share certain facilities. The Project will consist of photovoltaic (PV) solar modules, tracker components, power inverters, transformers, an electrical collection system, one or two project substations, a shared switchyard, battery storage, access roads, and a shared gen-tie line to the existing Southern California Edison (SCE) Red Bluff Substation.

## 2 SITE CHARACTERISTICS

### 2.1 Regional Setting

The Project site is in the central portion of Chuckwalla Valley, east of Palm Springs in the Colorado Desert. The elevation of Chuckwalla Valley ranges from less than 400 feet above mean sea level (amsl) at Ford Dry Lake to approximately 1,800 feet amsl west of Desert Center and along the upper portions of the alluvial fans that surround the valley perimeter. The surrounding mountains rise to over 3,000 feet amsl. The topography of the Project site generally slopes downward toward the northeast at gradient of less than 1 percent. Ground surface elevations at the Project site itself ranges from approximately 485 feet amsl in the northeast to 645 feet amsl in the southwest.

Anthropogenic features and land use near the Project site include agriculture, renewable energy, energy transmission, and historical military operations. Adjacent and nearby land uses are summarized in Table 1 and shown on Figure 1.

**Table 1.** Adjacent and Nearby Land Uses

Direction	LAND USES
NORTH	Active/fallow agriculture, Desert Lily Preserve, Joshua Tree National Park
SOUTH	Proposed Victory Pass Solar Project, transmission lines, active/fallow agriculture, I-10, Southern California Edison's Red Bluff substation
EAST	Athos Solar Project, Palen Solar Project, active/fallow agriculture, Palen Dry Lake ACEC
WEST	Multispecies linkage, Athos Solar Project, proposed Oberon Solar Project, transmission lines, Highway 177

## **2.2 Hydrology**

The Project site resides within the Colorado River Hydrologic Region (HR). The Colorado River HR covers approximately 13 million acres (20,000 square miles) in southeastern California and is the most arid HR in California with annual precipitation averaging 5.5 inches (DWR 1994). The Project site is in the Big Wash HUC 10 Hydrologic Areas, which flow to closed basins, not connected with the Colorado River or other traditional navigable waters (Figure 2). Palen Dry Lake and Ford Dry Lake represent the lowest elevations within the basin.

Desert washes within this region are almost always dry but contract and expand dramatically in size due to extreme variations in flows, which can range from high-discharge floods to extended periods when surface flow is absent. The Project site lies between the alluvial fans emanating from the Eagle Mountains to the west, Chuckwalla Mountains to the south, and Coxcomb Mountains to the north.

The Project site is situated in the lower alluvial fan that is characterized by less stabilized soils consisting of finer sand and silt, compared to the upper alluvial fan that supports more stabilized, rocky soils with well-defined channels. The topography the Project site is relatively flat with gradients of less than two percent.

Alluvial processes across the Project site generally flow from southwest to northeast. The I-10 (about 2 miles south of the Project site) crosses the alluvial fan that emanates from the Chuckwalla Mountains. The I-10 and associated wing dikes, which were constructed over 45 years ago, have altered natural surface flows from dozens of meandering small alluvial washes into concentrated discrete channels. Lancaster et al. (2014) noted that changes to drainage patterns resulting from the construction of I-10 translate into downstream hydrological degradation, focusing surface flow into freeway undercrossings and rendering portions of the alluvial fan less active than under historical conditions. Minor washes located in the hydrological shadow of I-10 were cut off from upstream flows and therefore transport lower volumes of water and entrained sediment. Major, culverted washes received more surface flow and distribute a higher volume and fine sediment compared to conditions that preceded the construction of I-10. These effects persist on the Project site under current conditions.

## **2.3 Soils**

Soils mapped on the Project site consist of two soil types per the United States General Soils Map. The southwestern portion of the site is mapped as the Vaiva-Quilotosa-Hyder-Cipriano-Cherioni map unit characterized by soils with high percentage (greater than 65 percent) of sand with moderate susceptibility to wind erosion. The northeastern portion of the site is mapped as Rositas-Dune-Land Carsitas. It is characterized by soils with a very high sand percentage (greater than 95 percent) and is highly susceptible to wind erosion (Figure 3).

## **2.4 Sand Transport System**

The Project site is located within the Chuckwalla Valley, a region of active aeolian (wind-blown) sand migration and deposition. Aeolian processes play a major role in the creation and establishment of sand dune formations and habitat in the Chuckwalla Valley and those within the Project site. Aeolian sands

(dunes, sand fields, and similar habitats) are important habitats for certain plants and animals, including Mojave fringe-toed lizard.

In conjunction with the DRECP process, the Department of Conservation's California Geological Survey prepared a regional Eolian System Mapping Report for Eastern Riverside County in 2014 (Lancaster et al. 2014; note that eolian and aeolian are alternate spellings of the same word).

Active aeolian sand transport areas were consistent with analysis conducted by Kenney et al (2014) with active aeolian sand migration in the northern and northeastern boundaries and with aeolian sand sources in the remainder of the site (Figure 4). The active areas of aeolian sand migration areas have finer sand that is suitable habitat for sensitive wildlife (such as Mojave fringe toed lizard) and plant species (such as Harwood's milkvetch). There is some instability over time and space, as sand corridors expand, contract or migrate with changing weather and climate. A more recent analysis by Kenney in 2017 shows that the remainder of the site, that has aeolian sand sources, has a low to moderate sand migration rate (Kenney 2017).

## 2.5 Rainfall

Measurements of precipitation during winter (October through March) and summer (April through September) periods are important in determining the efficacy of both wildlife and special status plant surveys. Data were obtained from the Western Regional Climate Center (WRCC 2020) for the most proximate stations to the Project site: Blythe Airport and Eagle Mountain weather stations (approximately 40 miles and 10 miles from the Project site, respectively). Historical rainfall data from 2009 to 2020 were totaled and averaged (Table 2). Over the period of analysis, the highest winter rainfall occurred in 2010 and highest summer rainfall occurred in 2012. Winter rains prior to the spring 2020 survey were above average over the last analysis period.

**Table 2.** Rainfall Summary

Year	Winter - October to March (inches)*	Summer - April to September (inches)*
2010	4.8	0.1
2011	2.5	1.2
2012	1	3.31
2013	1.5	2.6
2014	0.7	1.2
2015	2.1	1.3
2016	1.5	0.7
2017	3.4	1.1
2018	0.1	0.5
2019	2.6	0.165
2020	3.6	-
<b>Seasonal Average</b>	<b>2.2</b>	<b>1.2</b>

\*Source: Western Regional Climate Center (WRCC 2020): Blythe Airport and Eagle Mountain weather stations

## 2.6 Vegetation Communities

Vegetation communities in the Project site were mapped and classified by botanists, using Holland (1986) and cross-referencing with *A Manual of California Vegetation, 2nd edition* (Sawyer et al. 2009) and the National Vegetation Classification System (NVCS) referenced in the DRECP. Vegetation was mapped by drawing vegetation polygons on aerial images in the field. These field maps were then digitized into GIS shapefiles using ArcGIS Pro and one-foot pixel aerial imagery on a diagonal flat screen monitor at the office. The smallest mapping unit delineated was approximately 0.10 acres; most mapped vegetation boundaries are accurate to within approximately 10 feet.

The vegetation communities map (Figure 5) was generated from ArcGIS shapefiles; the shapefiles were used to calculate areas of each vegetation type and may be viewed at larger scale for management or analysis purposes, if needed. Any vegetation map is subject to imprecision for several reasons:

- Vegetation types tend to intergrade on the landscape so that there are no true boundaries in the vegetation itself. In these cases, a mapped boundary represents best professional judgment.
- Vegetation types as they are named and described tend to intergrade; that is, a given stand of real-world vegetation may not fit into any named type in the classification scheme used. Thus, a mapped and labeled polygon is given the best name available in the classification, but this name does not imply that the vegetation unambiguously matches its mapped name.
- Vegetation types tend to be patchy. Small patches of one named type are often included within mapped polygons of another type. The size of these patches varies, depending on the minimum mapping units and scale of available aerial imagery.

The majority of the Project site is creosote bush scrub. There are two primary natural vegetation communities (creosote bush scrub and desert dry wash woodland) as well as one distinct natural habitat type (desert pavement). One vegetation community (desert dry wash woodland/microphyll woodland) is identified by BLM (NECO Plan 2002 and DRECP 2016) and CDFW (2010) as sensitive due to the association with alluvial processes and would likely be considered California State jurisdictional waters. Vegetation communities on the Project site are shown on Figure 5.

### 2.6.1 Sonoran Creosote Bush Scrub

Sonoran creosote bush scrub has a State Rarity rank of S5 (CDFW 2020d), being demonstrably secure, and is not designated as a sensitive plant community by BLM. It is synonymous with *Larrea tridentata* - *Ambrosia dumosa* alliance (Sawyer et. al 2009) and *Lower Bajada and Fan Mojavean – Sonoran Desert Scrub* (NVCS). Sonoran creosote bush scrub occurs on well-drained, secondary soils of slopes, fans, and valleys and is the basic creosote bush scrub habitat of the Colorado Desert (Holland 1986). This vegetation community occurs through a majority of the Project and is dominated by creosote bush, burro bush, and has an understory of annual buckwheat (*Eriogonum* sp.) and *Cryptantha* species with sparse desert ironwood (*Olneya tesota*) and palo verde (*Parkinsonia florida*) near the southeastern boundary.



### 2.6.2 Desert Saltbush Scrub

Desert saltbush scrub has a state rarity rank of S4 (CDFW 2020), being demonstrably secure, and is not designated as a sensitive plant community by BLM. It is synonymous to *Atriplex canescens* (fourwing saltbush) shrubland alliance (NVCS). Holland (1986) associates this community with relict interior dunes. This vegetation community is dominated by fourwing saltbush with sparse creosote bush and *Cryptantha* species within the understory. This vegetation community is located on sandier soils near the northern and western boundaries of the Project site.

### 2.6.3 Desert Dry Wash Woodland

Desert dry wash woodland is a sensitive vegetation community recognized with a rarity rank of S4 (CDFW 2018d) and as a BLM sensitive vegetation community (NECO 2002, DRECP 2012). Desert dry wash woodland is characteristic of desert washes and is likely to be regulated by CDFW as jurisdictional state waters. This community is synonymous with blue palo verde (*Parkinsonia florida*) - ironwood (*Olneya tesota*) (microphyll) woodland alliance (Sawyer et. al 2009) and Sonoran - Coloradan Semi Desert Wash Woodland / Scrub (NVCS). Holland (1986) describes this community as an open to relatively densely covered, drought-deciduous, microphyll (small compound leaves) riparian scrub woodland, often supported by braided wash channels that change following every surface flow event. Within the Project site, this vegetation community is dominated by an open tree layer of ironwood, blue palo verde, and smoke tree (*Psoralea arguta*) of at least 2-3% cover. The understory is a modified creosote scrub with big galleta grass (*Hilaria rigida*) and occurs only in small amounts along the southwestern boundary.

## 3 DATA COLLECTION METHODS

### 3.1 Special Status Species Definition

Special status species are those that have been afforded special recognition by federal, state, or local resource agencies or organizations, are often of relatively limited distribution, and typically have unique habitat conditions, which also may be in decline. Special status criteria include:

- Officially listed or candidates for listing by California or the federal government as endangered, threatened, of special concern, or rare under the California Endangered Species Act (CESA) or the Federal Endangered Species Act (ESA);
- Plants or animals which meet the criteria for listing as rare or endangered, as described in Section 15380 of the California Environmental Quality Act (CEQA);
- BLM Sensitive Species designated by the BLM California State Director;
- USFWS Sensitive Species;
- Plants listed in the CNPS Inventory of Rare and Endangered Plants of California (CNPS 2020);
- Wildlife species identified by CDFW as Species of Special Concern (CNDDDB 2020);
- Plants or animals included in the CDFW lists of Special Plants or Special Animals (CNDDDB 2020);
- Considered special-status species in local or regional plans, policies, or regulations, such as the NECO Plan/EIS
- Protected under other statutes or regulations (e.g., Migratory Bird Treaty Act, Bald and Golden Eagle Protection Act, Native Plant Protection Act, etc.)

Based upon review of the literature, a list of special-status wildlife species with potential to occur in the vicinity of the Project site was compiled (Appendix B). All surveys were conducted per DRECP DFA Biological Conservation Management Action (CMA) requirements for each species within the timing recommended, including full-coverage desert tortoise surveys and 4-visit burrowing owl surveys. Any modifications are further explained within each individual sensitive species section below.

## **3.2 Wildlife Surveys**

Full coverage wildlife surveys were conducted during the following periods:

- Fall surveys, full-coverage 10-meter transect surveys: October 9-12, 2019
- Spring surveys, full coverage, 20-m transects with 50-m buffer: March 21-25, 2020
- Burrowing owl modified burrow surveys #2-4: May 5-8, 2020; June 17-18, 2020; July 14-15, 2020

Wildlife surveys in 2019 employed belt transects approximately 10 meters (32.8 feet) apart in order to provide 100 percent (full) coverage for the proposed solar facility. Wildlife surveys were repeated in 2020 at 20-meter belt transects in conjunction with plant surveys with a 150-meter buffer, consistent with 2012 burrowing owl protocol surveys (Figure 6).

Survey crews in the fall of 2019 consisted of experienced wildlife desert biologists with at least one botanist with each crew. Survey crews in spring 2020 surveys consisted of primarily experienced botanists with at least one wildlife biologist per crew. Surveys were conducted by walking linear transects and visually searching for live individuals or sign of any sensitive species. All holes detected that may be inhabited by sensitive species were carefully inspected for potential occupancy, or sign of recent use as burrows or burrow complexes. Special emphasis was placed on searching around the bases of shrubs and along the banks of shallow washes. Burrows were carefully examined and assigned to the wildlife species that may have inhabited them based on indicator signs within the burrow or near the mouth of the burrow.

During wildlife surveys, biologists recorded all wildlife species observed, regardless of conservation status. Common species were tallied at the end of each transect and recorded throughout each day by each crew. During the spring 2020 surveys, additional avian counts were completed in the mornings during surveys, until 10 a.m.

All locational information for special status species observations and sign detected were recorded on digital Zerion iForms for any new data collected. The data was then uploaded to Collector as reference for Fall 2020 surveys to ensure that duplicate data was not taken. A handheld Global Positioning System (GPS) unit was used to collect backup data for each occurrence recorded with a unique identifier number that was also recorded on the digital Zerion iForms.

Incidental observations of all wildlife, regardless of conservation status, was also recorded for both 2019 and 2020 surveys (Appendix E).

### **3.2.1 Desert Tortoise**

Wildlife surveys conducted in fall 2019 conformed to full coverage desert tortoise protocol surveys (USFWS 2019a). All tortoise sign [e.g., live tortoises (all age classes), shell/bone/scutes, scats,

burrows/pallets, tracks, eggshell fragments, and courtship rings] observed was recorded within the known action area of the Project site. Incidental observations of desert tortoise sign were recorded in 2020 if they were not previously recorded. The condition of burrows, scat, and carcasses were categorized per the following class designations (USFWS 2009):

- Burrows:
  1. currently active, with desert tortoise or recent desert tortoise sign
  2. good condition (no evidence of recent use) - definitely desert tortoise
  3. deteriorated condition (including collapsed burrows) - definitely desert tortoise
  4. good condition - possibly desert tortoise
  5. deteriorated condition (including collapsed burrows) - possibly desert tortoise.
- Scat:
  1. wet (not from rain or dew) or freshly dried, obvious odor
  2. dried, with glaze, some odor, dark brown
  3. dried, no glaze or odor, signs of bleaching (light brown), tightly packed material
  4. dried, light brown to pale yellow, loose material, scaly appearance
  5. bleached, or consisting only of plant fiber
- Carcasses:
  1. < 1 year, fresh putrid, scutes mostly adhered, sheen on exposed scutes, unexposed bone waxy and solid;
  2. 1-2 years, scutes mostly adhered to bone, exposed scutes pale without sheen, unexposed bone silky;
  3. 2-3 years, scutes peeling off bone, unexposed scutes pale and without sheen, no growth ring peeling
  4. 4 years, shell bone falling apart, growth rings on scutes peeling; bone fissured
  5. > 4 years, disarticulated and scattered

### **3.2.2 Mojave Fringe-toed Lizard**

There is no protocol for surveying Mojave fringe-toed lizards, but during wildlife surveys, special attention was given to the search for live individuals in soft, sandier soils where the potential for the species to occur is high. In areas with a higher density of Mojave fringe-toed lizards observed within close proximity of one another (within 20 meters), groups of lizards were tallied and represented by a single data point on Project figures.

### **3.2.3 Couch's Spadefoot Toad**

A reconnaissance level survey for potential Couch's spadefoot toad habitat was conducted in conjunction with 2019 fall surveys searching for areas that may provide suitable habitat for reproduction. Areas where water may accumulate and retain for at least 2 weeks following heavy rain were recorded as potential Couch's spadefoot toad reproductive habitat to be inspected again following heavy rains during appropriate warmer temperatures for any sign of Couch's spadefoot toad.

### **3.2.4 Avian Species**

#### **3.2.3.1 Western Burrowing Owl**

Survey recommendations in both the 1993 California Burrowing Owl Consortium (CBOC) Guidelines and 2012 CDFW Staff Report include baseline data collection and an assessment of site use by burrowing owl. One full-coverage survey was conducted during the breeding season during 2020 spring surveys which was consistent with Phase II of the CBOC 1993 Guidelines and partially consistent with the 2012 CDFW Staff Report. Occupancy of burrowing owl habitat is confirmed at a site when at least one burrowing owl, or its sign at or near a burrow entrance, is observed within the last three years (CDFW 2012; CBOC 1993).

The first burrowing owl survey provided a greater level of coverage than the 30-meter spacing recommended in the 1993 CBOC Guidelines and the 20-meter spacing recommended in the 2012 CDFW Staff Report. This first survey was completed in conjunction with spring plant surveys. All burrows detected during this survey were assessed for wildlife occupancy, to ensure detection of any special status species, including burrowing owl, that may have occupied a burrow. The 20-meter transect spacing also increased the likelihood of flushing live burrowing owls during the survey. All sign of burrowing owl, including individuals, feathers, tracks, whitewash, pellets, and suitable burrows were recorded if present. An additional 50-meters of buffer around the Project site was also surveyed following the 2012 protocol survey.

A modification of the protocol 2012 survey recommendations was completed for the subsequent three surveys. The subsequent three surveys were modified as burrow inspections for all previously detected burrows, including mammal, potential tortoise, or burrowing owl burrows, with meandering pedestrian transects throughout the Project site where previous burrows were detected. All burrows were re-visited to check for any change in burrowing owl sign and were included as new burrowing owl sign if detected. Any new burrows observed during these burrow checks were added to the next check. These burrow checks were spaced at the same time intervals as the 2012 recommendations with at least 3 weeks of time passing between each session of burrow surveys.

#### **3.2.3.2 Golden Eagle**

Targeted surveys for golden eagles were not performed for the Project due to numerous surveys conducted in the Project vicinity and Chuckwalla Valley within the last ten years. A compilation of survey methodology and results from other projects that have conducted these surveys in the last ten years is provided in the results section of this report.

#### **3.2.3.3 Avian Counts**

Avian counts were conducted during spring 2020 surveys. Each survey team consisted of at least one avian biologist who was exclusively tasked with tallying all avian observations while walking with each survey team in the morning from the start of the survey until about 10:00 am, or earlier if weather conditions were unfavorable for avian detection (i.e. high wind). After these avian counts, the avian biologist would continue to note any incidental wildlife species observed while also continuing to help with any survey that was occurring.

### **3.2.5 Special Status Bat Species**

Targeted surveys for bats were not conducted and incidental observations of bats or bat roosts were not detected during wildlife surveys. Acoustic bat surveys previously conducted for a nearby project currently in construction, Palen Solar Energy Project, provides supplementary information about bat populations within the project vicinity. This is further discussed in the results section of this report.

### **3.2.6 Other Special Status Wildlife Species**

All sign of desert kit fox and American badger was recorded including live or dead individuals, scat, tracks, burrows, and burrow complexes. All burrows and burrow complexes were mapped and attributed, if possible, to species. If a burrow could not be attributed to species, it was recorded as a “canid” burrow, which may include desert kit fox, coyote, or domestic dog. Species usage for each burrow or complex was determined by the burrow size, shape, and sign (i.e. type of scat or scratches). If fresh tracks, scratches, freshly dug dirt or scat were found at a burrow or complex, it was categorized as active. The presence of old scat without tracks would indicate that a burrow or complex was inactive.

### **3.2.7 Wildlife Cameras**

Wildlife cameras were placed at seven locations near I-10 underpasses to determine wildlife movement in those areas – at least six camera locations were within the multispecies linkage. Cameras were secured to trees near the underpasses or hidden on wash edges among creosote bushes to deter theft. Cameras were set on high sensitivity, triggered by any movement to take photos in the daytime and evening with infrared light. Cameras were setup for one month in the spring of 2020. While cameras were in use, they were inspected weekly to ensure security and verify that battery levels and memory storage for cameras were sufficient.

## **3.3 Special Status Plants**

Focused special status plant surveys were conducted during the following periods:

- Fall surveys: full coverage, 10-meter transect surveys – October 9-12, 2019
- Spring Surveys: full coverage, 20-meter transect surveys with 50-meter buffer; March 21-25, 2020

Survey methodology was consistent with the following guiding documents:

- Guidelines for Conducting and Reporting Botanical Inventories for Federally Listed, Proposed, and Candidate Plants (USFWS 2000)
- Guidelines for Assessing the Effects of Proposed Projects on Rare, Threatened, and Endangered Plants and Natural Communities (CDFG 2009)
- CNPS Botanical Survey Guidelines (CNPS 2001)
- Survey Protocols for Survey and Manage Strategy 2: Vascular Plants (Whiteaker 1998)
- Survey Protocols Required for NEPA/ESA Compliance for BLM Special Status Plant Species (BLM 2009).

Based upon review of the literature, a list of special-status plant species with potential to occur in the vicinity of the Project site was compiled (Appendix C).

Plant surveys performed in fall of 2019 included visual coverage across the entire survey area in conjunction with wildlife surveys. Surveys employed belt transects approximately 10 meters apart in areas with native vegetation cover, access roads, and the gen-tie routes to provide 100 percent coverage in those areas. All surveyors were trained on diagnostic features and habitat notes special-status species that may occur and each crew of surveyors had at least one highly experienced botanist.

A focused plant survey was performed in spring 2020 that included visual coverage across the entire Project site. Surveyors employed belt transects spaced at approximately 20-meters apart. These surveys were completed in conjunction with avian counts in the mornings. Avian biologists chosen to support these botanical surveys were also skilled in plant identification and could easily integrate with the plant survey after completing avian counts in the morning.

Prior to beginning plant surveys in both the fall and spring, reference populations of sensitive plants were visited to ensure that timing for surveys was sufficient and that most sensitive plant species would be identifiable (Table 3).

**Table 3.** Plant Reference Population Visits

Date	Plant Species	Location	Status
10/9/2019	Crucifixion thorn ( <i>Castela emoryi</i> )	Desert Center	Observed
10/9/2019	Desert unicorn plant ( <i>Proboscidea althaeifolia</i> )	Desert Center	Observed
1/26/2020	Harwood's eriastrum ( <i>Eriastrum harwoodii</i> )	Desert Lily Sanctuary, Hwy 62/Ironage Rd	Observed, but not yet flowering – estimated to begin flowering in 3 weeks
1/26/2020	Harwood's milkvetch ( <i>Astragalus harwoodii</i> )	Desert Lily Sanctuary, Hwy 62/Ironage Rd	Too early to differentiate between other <i>Astragalus</i> sp.
1/26/2020	Ribbed cryptantha ( <i>Cryptantha costata</i> )	Desert Lily Sanctuary, Hwy 62/Ironage Rd	Too early to differentiate from other <i>Cryptantha</i> sp.
1/26/2020	<i>Ditaxis serrata</i> var. <i>californica</i>	Desert Center	Could not find even after contacting original specimen collector in area
3/17/2020	Harwood's eriastrum ( <i>Eriastrum harwoodii</i> )	Desert Lily Sanctuary, Desert Center	Observed flowering
3/17/2020	Harwood's milkvetch ( <i>Astragalus harwoodii</i> )	Desert Lily Sanctuary, Desert Center	Observed flowering
3/17/2020	Ribbed cryptantha ( <i>Cryptantha costata</i> )	Near Palen Solar Project	Observed

During plant surveys, botanists recorded all plant species, regardless of conservation status. All locational information for special status species observations were recorded on digital Zerion iForms for any new data collected. The data was then uploaded to Collector as reference for spring 2020 surveys to ensure that duplicate data from the previous fall season was not taken. A handheld Global Positioning System (GPS) unit was used to collect backup data for each occurrence recorded with a unique identifier number that was also recorded on the digital Zerion iForms. All necessary data submissions have been submitted from electronic data collections as required per protocols listed.

**Table 4.** Special-status Wildlife and Plant Survey Personnel and Dates \*

Date Ranges	Survey Type	Surveyors
10/9/2019-10/12/2019	Fall 2019 10-m surveys, wildlife and plant	M. Adams, T. Alvey, M. Baker, M. Bassett, K. Black, M. Cloud-Hughes, E. Bowen, M. Bratton, S. Clegg, L. Chow, M. Dipane, M. Honer, S. Hoss, T. Hobbs, C. Keaton, D. Kesonie, M. Lavender, M. Lopez, A. Mach, W. McBride, M. Moon, S. Nielsen, S. Nelson, B. Payne, B. Sandstrom, C. Slaughter, E. Thorn, J. St Pierre, M. Sally, R. Woodard, Z. Webb, M. Wegmann, J. Yerger, M. Zhuo
3/21/2020-3/25/2020	Spring 2020 20-m plant surveys/BUOW survey #1	M. Adams, M. Baker, K. Black, L. Chow, M. Honer, M. Cloud-Hughes, D. Kesonie, W. McBride, M. Lavender, M. Lopez, S. Nielson, J. St Pierre, J. Yerger
5/1/2020-5/29/2020	Spring 2020 wildlife cameras at underpasses	L. Chow, Z. Webb, M. Lavender
5/6/2020-5/8/2020	BUOW survey #2	Z. Webb, L. Chow
5/20/2020-5/21/2020	Jurisdictional delineations**	L. Chow, E. Thorn, D. Kesonie, J. St Pierre, S. Nielson
6/17/2020-6/18/2020	BUOW survey #3	A. Schaub, M. Lavender
7/14/2020-7/15/2020	BUOW survey #4	Z. Webb, M. Lavender

\* daily survey efforts are detailed in Table A-1 of Appendix A

\*\*Jurisdictional delineation methods and results are provided in a separate jurisdictional waters report for the Project

## 4 RESULTS

### 4.1 Special Status Wildlife

Sixty-one special status wildlife species were reviewed for their potential to occur within the Project site and its vicinity using information gathered from regional plans and database records (Appendix B). Several species were determined to have a low probability of occurrence due to the absence of suitable habitat. Special status wildlife species observed within the Project site or with moderate potential to occur based on the presence of suitable habitat are discussed further in this section. Appendix A contains tables of all survey results. A comprehensive list of wildlife species observed during previous surveys is included in Appendix D. Conservation status for wildlife species is defined below:

- Federal
  - FE = Federally listed endangered: species in danger of extinction throughout a significant portion of its range
  - FT = Federally listed, threatened: species likely to become endangered within the foreseeable future
  - FCT = Proposed for federal listing as a threatened species
  - BCC = Fish and Wildlife Service: Birds of Conservation Concern
  - BLMS = BLM Sensitive
- State
  - SSC = State Species of Special Concern
  - CFP = California Fully Protected
  - SE = State listed as endangered
  - ST = State listed as threatened
  - WL = State watch list
  - CPF = California Protected Furbearing Mammal
  - CPGS = California Protected Game Species

#### 4.1.1 Mojave Desert Tortoise: ST, FT

##### Background

Mojave desert tortoises (*Gopherus agassizii*) live north and west of the Colorado River in the Mojave Desert of California, southern Nevada, northwestern Arizona, and southwestern Utah, and in the Sonoran (Colorado) Desert in southern California (USFWS 1990). Desert tortoises inhabit a variety of habitats from flats and slopes dominated by creosote bush – white bursage communities, where a diversity of perennial plants is relatively high, to a variety of habitats in higher elevations. Tortoises are found most often on gentle slopes with sandy-gravel soils. Soils must be appropriately soft for digging burrows, but firm enough so that burrows do not collapse (Anderson et al., 2000). Tortoises typically prefer habitats with abundant annual forbs, grasses and cactus, which constitute its primary food sources. Plant species that have high potential for potassium excretion (high-PEP) may be critical to the diet of desert tortoise (Oftedal 2002; Oftedal et. al 2002).

The Project site is located within the Colorado Desert Recovery Unit for Mojave desert tortoise (USFWS 2011) and is less than 2 miles north of USFWS-designated critical habitat for desert tortoises, which is also designated as a Desert Tortoise Conservation Area (TCA) in the DRECP (Figures 1,7, and 8). The 2019 density (/km<sup>2</sup>) of tortoises with a midline carapace length of greater than or equal to 180 mm within the Colorado Desert Recovery Unit strata are as follows: Chuckwalla 1.8 tortoises per km<sup>2</sup> (less than two miles to the south of the project site), Joshua Tree 3.1 per km<sup>2</sup> (approximately five miles north of the project site), Pinto Mountains 1.7 per km<sup>2</sup>, approximately 25 miles northwest from the Project site), Chocolate Mountain 7 per km<sup>2</sup> approximately 20 miles south of the Project site, and Fenner 2.8 per km<sup>2</sup>, approximately 70 miles north of the Project site (USFWS 2020). Surveys in the Chemehuevi stratum were not conducted in 2019. In 2018 the density of tortoises within the Chemehuevi stratum was 2.9 tortoises per km<sup>2</sup> (approximately 60 miles from the project site) (USFWS 2019b).

Predicted Mojave desert tortoise occupancy values of 0.3 or above are appropriate for identifying suitable habitat in this low desert region (BLM 2012). Desert tortoise habitat has the lowest predicted occupancy levels in the northernmost portion of the Project site (less than 0.3) and increases as it moves southwest. The highest occupancy levels of 0.5-0.6 is in the southwest portion of the site (Nussear et al. 2009). These predicted occupancy values do not account for habitat degradation resulting from existing anthropogenic features (Nussear et al. 2009), which would further reduce the occurrence probability in disturbed areas. Desert tortoise habitat connectivity is discussed in Section 4.2, Wildlife Movement.

Since no live Mojave desert tortoises were documented during protocol level surveys for the species, the USFWS population estimate methodology cannot be used to calculate the population density. There was minimal desert tortoise sign on the Project site. The only desert tortoise sign observed during field surveys on the Project site was a class 4 desert tortoise burrow on the western boundary of the Project site. This is consistent with the predicted occupancy model, with the only observed sign occurring in the area with occupancy values of 0.4-0.5 (Figure 7, Appendix A-2).



#### **4.1.2 Mojave fringe-toed lizard: SSC, BLMS**

The Mojave fringe-toed lizard (*Uma scoparia*) occupies arid, sandy, sparsely vegetated habitats and is associated with creosote scrub throughout much of its range (Jennings and Hayes 1994). It is found within and around aeolian sand habitats in the deserts of Los Angeles, Riverside, and San Bernardino Counties in California and La Paz County in Arizona (Hollingsworth and Beaman 1999; Stebbins 1985; Murphy et al. 2006). Within these regions, it occurs at more than 35 sand dune complexes in California and one in Arizona (Jarvis 2009). Nearly all records for this species are associated with present-day and historical drainages and sand dune complexes associated with three major river systems with blow sand: Amargosa River, Mojave River, and Colorado River (BLM 2015).

Mojave fringe-toed lizards normally hibernate from November to February, emerging from hibernation sites from March to April. The breeding season is April to July (Mayhew 1965). From May to September, they are active in mornings and late afternoon, but seek cover during the hottest parts of the day. They burrow in the sand for both cover from predators and protection from undesirable temperatures (Stebbins 2003), though they also will seek shelter in rodent burrows.

As this species requires loose, wind-blown sand, its distribution within the survey areas is consistent with the presence of suitable soil conditions. All detections for Mojave fringe-toed lizard were concentrated within areas where the DRECP distribution model for the species overlapped with the more active sand transport areas. Observations were primarily on the sandiest parts of the site with a total of 32 areas of observations within the Project site on the northern and eastern most boundaries (Figure 7, Appendix A-2).

#### **4.1.3 Couch's Spadefoot Toad: SSC, BLMS**

Couch's spadefoot toad (*Scaphiopus couchii*) is often found in shortgrass plains, mesquite savannah, creosote bush desert, thorn forest, and tropical deciduous forest (Mexico) and other areas of low rainfall (Stebbins 2003). It is considered an opportunistic species because it only appears when rainfall forms temporary pools and potholes with water lasting longer than 10-12 days, which are required for breeding, hatching, and metamorphosis. Runoff basins at the base of sand dunes are also sites of reproduction (Mayhew 1965). In California, it is known to occur in the low desert region, especially the Colorado River corridor. It burrows underground or occupies rodent burrows when inactive.

Couch's spadefoot toad was not observed, but suitable breeding habitat may be present within the Project site in areas where water accumulation may occur. Nine areas were identified during fall 2019 surveys as potential breeding habitat (Figure 7, Appendix A-2) where water may accumulate after rainfall but sufficient rainfall in warmer temperatures is yet to occur to determine whether they hold enough water for breeding or any occupancy of the species. It is expected to be a low potential of occurrence for the species since there have been no records of the species at nearby projects.

#### **4.1.4 Western Burrowing Owl: SSC, BCC, BLMS**

The Western burrowing owl (*Athene cunicularia hypugaea*) inhabits arid lands throughout much of the western United States and southern interior of western Canada (Haug et al. 1993). Suitable habitat for western burrowing owl includes open habitat with available burrowing opportunities, including

agricultural fields (active and fallow), creosote scrub, desert saltbush, ephemeral washes, and ruderal areas.

Burrowing owls are unique among the North American owls in that they nest and roost in abandoned burrows, especially those created by ground squirrels, kit fox, desert tortoise, and other wildlife. Burrowing owls have a strong affinity for previously occupied nesting and wintering sites and will often return to previously used burrows, particularly if they had successful reproduction in previous years (Gervais et al. 2008). They generally depend on other species to dig suitable burrows for use but may also use anthropogenic surrogate burrows such as rubble piles or drainage pipes. If formerly occupied burrows are badly damaged or collapsed, burrowing owls cannot repair them and must seek alternate sites. The southern California breeding season (defined as the time from pair bonding of adults to fledging of the offspring) generally occurs from February to August, with peak breeding activity from April through July (Haug et al. 1993).

In the Colorado Desert, burrowing owls generally occur at low densities in scattered locations, but they can be found in much higher densities near agricultural lands where rodent and insect prey tend to be more abundant (Gervais et al. 2008). Burrowing owls tend to be opportunistic feeders, and a large portion of their diet consists of beetles, grasshoppers, and other larger arthropods. The consumption of insects increases during the breeding season (Haug et al. 1993). Small mammals, especially mice and voles (*Microtus* and *Peromyscus* spp.) are important food items, and other prey animals include herpetofauna, young cottontail rabbits, bats, and birds such as sparrows and horned larks.

Five observations of live individuals flying/perching, eight observations of live individuals at burrows with sign, and five burrows with sign (whitewash, pellets, or feathers) were observed (Figure 9, Appendix A-3).

#### **4.1.5 Golden Eagle: CFP, WL, BCC, BLMS**

##### **Background**

Golden eagles are typically year-round residents throughout most of their western United States range. They breed from late January through August with peak activity occurring from March through July (Kochert et al. 2002). Habitat for golden eagles typically includes rolling foothills, mountain areas, and deserts. Golden eagles need open terrain for hunting and prefer grasslands, deserts, savanna, and early successional stages of forest and shrub habitats. Golden eagles primarily prey on rabbits and rodents but will also take other mammals, birds, reptiles, and some carrion (Kochert et al. 2002). They generally nest in rugged, open habitats with canyons and escarpments, often with overhanging ledges and cliffs or large trees used as cover.

Recent data analysis and population modeling suggest the status of the golden eagle population in the western United States is gradually declining towards an equilibrium of about 26,000 individuals, down from an estimated 34,000 in 2009 and 2014 (USFWS 2016). The future population estimate relies on the continuation of current ecological and biological conditions. It was estimated that 3,400 golden eagles die annually from anthropogenic causes in the United States and suggest a level of sustainable take is

approximately 2,000 individuals annually (USFWS 2016). Additional unmitigated mortality will steepen the rate of decline that the golden eagle population is presently undergoing (USFWS 2016).

The Project site does not have any suitable nesting habitat for golden eagles, but there is suitable foraging habitat. The nearest known cliff nest site that have the potential for golden eagle use, is approximately 4 miles from the Project site. No golden eagles were observed during surveys on the Project site.

#### Regional Surveys

No focused golden eagle surveys were conducted specifically for this Project. No live golden eagles were observed flying over the Project site during the field surveys described above (Section 3.2.4). Golden eagle surveys and raptor surveys have been conducted on a multitude of projects within 10 miles of the Project vicinity since 2010. The most recent survey was conducted in spring 2020 – general location information to share this information is pending approval from USFWS. Type of survey and results for regional golden eagle surveys between the years of 2010-2020 are summarized in Table 5 below. The highest concentration of overlapping surveys occurs within the Project vicinity (Figure 10).

**Table 5.** Regional Golden Eagle Surveys

Year	Type of Survey	Associated Project (s)	Surveying Firm	Golden Eagle Observations
2010	Aerial survey	Desert Sunlight Solar Project, Genesis Solar Project, Palen Solar Project	Wildlife Research Institute	1 active nest in Coxcomb Mountains, 1 active territory in Eagle Mountains
2011	Aerial eagle (not nesting) and transect survey	Other research survey	West	No observations in area surveyed
2011	Aerial and ground	Regional Nest Survey	BioResource Consultant	No observations in area surveyed
2011	Aerial survey	Joshua Tree National Park	Wildlife Research Institute	4 territories active - Eagle Mountains-West Central, Eagle Mountains - West Northwest, Hexie Mountains - Central, Little San Bernardino - East); the Eagle Mountain territories were productive - had a total of 3 young observed
2011	Ground survey	Desert Harvest Solar Project	Bloom Biological	No active nests, 1 golden eagle sighting
2012	Aerial (not nesting) and transect survey, tracking eagles	Other research survey	West and Duerr et al	No observations in area surveyed
2012	Ground survey	Desert Sunlight Solar Project	Ironwood Consulting	No active nests - 7 golden eagle sightings (6 in Eagle Mountains, 1 in Coxcomb Mountains)
2013	Tracking eagles	Other research survey	West and Duerr et al	No observations in area surveyed

Year	Type of Survey	Associated Project (s)	Surveying Firm	Golden Eagle Observations
2013	Ground survey	BLM raptor-raven nest survey	Corvus Ecological	No observations in area surveyed
2013	Ground survey	Desert Sunlight Solar Project	Corvus Ecological	No active nests, 4 golden eagle sightings
2013	Air and ground survey, camera traps	Palen Solar Project	Bloom Biological	1 subadult at bait station during all 5 weeks; 3rd year flying along cliffs
2014	Air and ground survey	BLM raptor-raven nest survey	Boarman	No observations in area surveyed
2015	Ground survey	BLM raptor-raven nest survey	Corvus Ecological	No observations in area surveyed
2020	Ground survey	BLM raptor-raven nest survey	Corvus Ecological	3 nests in Joshua Tree National Park (general locational information pending)
2020	Variable radius point count	Chuckwalla CHU	Corvus Ecological	General locational information pending

#### 4.1.6 Le Conte's Thrasher: SSC

In California, Le Conte's thrasher (*Toxostoma lecontei*) is a resident in the San Joaquin Valley and the Mojave and Colorado Deserts (Weigand and Fitton 2008). This pale gray bird occurs in desert flats, washes and alluvial fans with sandy and/or alkaline soil and scattered shrubs. Preferred nest substrate includes thorny shrubs and small desert trees and nesting rarely occurs in monotypic creosote scrub habitat or Sonoran Desert woodlands (Prescott 2005). Breeding activity occurs from January to early June, with a peak from mid- March to mid-April. Le Conte's thrashers forage for food by digging and probing in the soil. They eat arthropods, small lizards and snakes, and seeds and fruit; the bulk of their diet consists of beetles, caterpillars, scorpions, and spiders. Suitable foraging habitat occurs throughout the site, and suitable nesting habitat is present only in the desert dry wash woodland areas (a small area in the southwest corner of the site; see Figure 5), so the potential for the species to nest on the Project site is low due to the small area of suitable nesting habitat. One live individual was observed and heard singing (Figure 9, Appendix A-3).

#### 4.1.7 Prairie Falcon: WL, BCC

The prairie falcon (*Falco mexicanus*) is on the CDFW watch list and is a USFWS Bird of Conservation Concern. It inhabits dry environments in the North American west from southern Canada to central Mexico. It is found in open habitat at all elevations up to 3,350 m, but is associated primarily with perennial grasslands, savannahs, rangeland, some agricultural fields, and desert scrub areas. Prairie falcons require cliffs or bluffs for nesting though will sometimes nest in trees, on power line structures, on buildings, or inside caves or stone quarries. Ground squirrels and horned larks are the primary food source, but prairie falcons will also prey on lizards, other small birds, and small rodents (Zeiner 1990).

There were seven observations of prairie falcon in flight throughout the Project site – many observations were from the same day from different survey crews, so it was likely repeat observations of the same individual. The entire Project site contains suitable foraging habitat for this species but does not have suitable nesting habitat (Figure 9, Appendix A-3).

#### **4.1.8 Loggerhead Shrike: SSC (nesting), BCC**

Loggerhead shrikes (*Lanius ludovicianus*) are small predatory birds that are common year-round residents throughout most of the southern portion of their range, including southern California. In southern California, they are generally much more common in interior desert regions than along the coast (Humple 2008). They can be found within lowland, open habitat types, including creosote scrub and other desert habitats, sage scrub, non-native grasslands, chaparral, riparian, croplands, and areas characterized by open scattered trees and shrubs. Loss of habitat to agriculture, development, and invasive species is a major threat; this species has shown a significant decline in the Sonoran Desert (Humple 2008). Loggerhead shrikes initiate their breeding season in February and may raise a second brood as late as July; they often re-nest if their first nest fails or to raise a second brood (Yosef 1996). In general, loggerhead shrikes prey upon large insects, small birds, amphibians, reptiles, and small rodents over open ground within areas of short vegetation, usually impaling prey on thorns, wire barbs, or sharp twigs to cache for later feeding (Yosef 1996). Suitable habitat for loggerhead shrike is found throughout the Project site. There were 10 observations of live individuals flying and perching during 2019-2020 surveys (Figure 9, Appendix A-3).

#### **4.1.9 Gila Woodpecker: CE, BLMS, BCC**

Gila woodpecker is predominantly a permanent resident across its range in areas of southeast California, southern Nevada, central Arizona, extreme southwest New Mexico, and parts of Mexico. The Gila woodpecker is an uncommon to fairly common resident in Southern California along the Colorado River, and locally near Brawley, Imperial County (Garrett and Dunn 1981). A pair of Gila woodpeckers was incidentally observed feeding young near the Corn Springs Campground, approximately 6 miles southwest of the Project site (Ironwood 2019). Suitable habitats include riparian woodlands, uplands with concentrations of large columnar cacti, old-growth xeric-riparian wash woodlands, and urban or suburban residential areas (Rosenberg et al. 1987; Edwards and Schnell 2000). Gila woodpeckers prefer large patches of woody riparian vegetation for nesting (greater than 49 acres), but they have also been documented in various habitat types, such as desert washes (McCreedy 2008) and residential areas (Mills et al. 1989). They excavate cavity nests in large riparian trees such as cottonwoods.

In California, their primary habitat is cottonwood-willow riparian woodland. Where Gila woodpeckers occur in dry desert wash woodlands, they excavate cavity nests in large blue palo verdes (McCreedy 2008). They also may nest in ornamental trees including palms. Availability of suitable nesting trees is a limiting factor in breeding habitat suitability (Grinnell and Miller 1944). Potentially suitable habitat within the Project site is found in desert washes in palo verde or ironwood trees large enough for cavity nests. The probability of this species nesting on the Project site is low since desert dry wash woodland and larger trees on the Project site is sparse (see Figure 5). Where Gila woodpeckers occur, they generally are loud and conspicuous, and readily located by field biologists. No Gila woodpeckers were observed during the field surveys.

#### **4.1.10 California Horned Lark: WL**

The California horned lark (*Eremophila alpestris actia*) is found throughout California except the north coast and is less common in mountainous areas. It prefers open areas that are barren or with short vegetation including deserts, brushy flats, and agricultural areas, and includes creosote scrub. Eggs are

laid March to early June, and it frequently lays a second clutch (Zeiner 1990). There are numerous records in western Riverside County (CNDDDB 2020). Suitable habitat for foraging and nesting is found throughout the Project site and California horned larks were observed frequently during the surveys. Observation locations were not mapped because of the low conservation status (WL) and widespread occurrence throughout the site.

#### **4.1.11 Black-tailed Gnatcatcher: WL**

Black-tailed gnatcatchers (*Polioptila melanura*) are permanent residents from southeastern California and Arizona to southern Texas and northern Mexico. They are found in arid scrublands, desert brush, and dry washes amongst creosote bush, ocotillo, mesquite, paloverdes, and cactus. They live in pairs all year-round, defend their territory, and forage for small insects amongst low shrubs and trees. The Project site contains suitable foraging and potential nesting habitat for this species throughout the site and there were several incidental observations during surveys and avian counts. Observation locations were not mapped because of the low conservation status (WL) and widespread occurrence throughout the site.

#### **4.1.12 Sonora Yellow Warbler: SSC, BCC**

The Sonora yellow warbler (*Setophaga petechia sonorana*) occurs principally as a migrant and summer resident from late March through early October, and breeds from April to late July (Dunn and Garrett 1997). The Sonora yellow warbler breeds only along the lower Colorado River in California, and from southern Arizona and southwest New Mexico to north-central Mexico and possibly the Colorado River Delta. It breeds in early April and nests mainly from mid-May through July (Rosenberg et al. 1991). During breeding season, it generally nests and forages in riparian shrubs and trees close to water. Its diet includes ants, bees, wasps, caterpillars, beetles, true bugs, flies, and spiders (Beal 1907, Shuford 2008). The Project site supports suitable foraging habitat during migration in the desert dry wash woodland areas, but there is no suitable nesting habitat present on the site. There were no observations of the Sonora yellow warbler on the Project site.

#### **4.1.13 Short Eared Owl: SSC**

The short-eared owl (*Asio flammeus*) is a widespread winter migrant in central and western California, and generally present from September through April. It is an uncommon winter migrant in southern California. Habitat requirements include grasslands, prairies, dunes, meadows, irrigated lands, and wetlands. Short-eared owls generally require dense vegetation for roosting and nesting (Shuford 2008). The Project site does not support suitable nesting habitat for short eared owl due to the sparse vegetation. However, the species may be found incidentally during migration while foraging near irrigated areas in the adjacent fish farms and residences. One individual was observed flying and roosting on the Project site (Figure 9, Appendix A-3).

#### **4.1.14 Ferruginous Hawk: WL, BCC**

The ferruginous hawk (*Buteo regalis*) is an uncommon winter resident and migrant at lower elevations and open grasslands in the Central Valley and Coast Ranges, and a fairly common winter resident of grasslands and agricultural areas in southwestern California (Garrett and Dunn 1981). This species frequents open grasslands, sagebrush flats, and desert scrub. Prey items include lagomorphs, small

mammals, reptiles and amphibians (Zeiner 1990). There is potential foraging habitat throughout the Project site that ferruginous hawks could use during wintering or migration seasons. The site is outside the Ferruginous hawk's breeding range and is not expected in the area during nesting season. There were no observations of ferruginous hawk on the Project site.

#### **4.1.15 Swainson's Hawk: ST, BBC**

Swainson's hawk (*Buteo swainsoni*) breeds in open habitats throughout much of the western United States and Canada, and in northern Mexico. In California, breeding populations of Swainson's hawks occur in desert, shrub and grasslands, and agricultural habitats with tree rows; however, most of the state's breeding sites are in the Great Basin and Central Valley (Woodbridge 1998). The only desert breeding occurrences are in the Antelope Valley, over 200 miles northwest of the Project site. These birds favor open habitats for foraging, and are near-exclusive insectivores as adults, but may also forage on small mammals and reptiles. The project site provides potential migration season foraging habitat but is well outside the nesting range. Swainson's hawk may be found throughout the project site during migration and one individual was observed in flight during surveys (Figure 9, Appendix A-3).

#### **4.1.16 American Peregrine Falcon: FP, BCC**

The American peregrine falcon (*Falco peregrinus anatum*) was formerly listed under CESA and ESA but has been delisted under both Acts. In California, its range is primarily central to northern California, with wintering habitat and (more recently) nesting occurrences located in southern California. Migrants occur along the coast and in the western Sierra Nevada in spring and fall. It breeds mostly in woodland, forest, and coastal habitats, and favors open landscapes with cliffs as nest sites. They are found irregularly in the southern desert region, generally during migratory and winter seasons but also during breeding season in recent years. They nested historically in desert mountain ranges near the Colorado River (Rosenberg et al. 1991; Patten et al. 2003) and may be re-occupying this historical part of their nesting range as their populations recover. Their diet consists primarily of birds and bats (Zeiner 1990). Waterfowl and shorebirds make up a large proportion of their prey, and nest sites are often within foraging range of large water bodies. Suitable migratory or foraging habitat is present throughout the Project site, but no suitable nesting habitat is present. There were no observations of the American peregrine falcon on the Project site.

#### **4.1.17 Vaux's Swift: SSC**

Vaux's swift (*Chaetura vauxi*) is a summer resident of northern California and a fairly common migrant throughout most of the state in spring and fall. It roosts in hollow trees and snags, and often in large flocks. Vaux's swifts feed exclusively on flying insects (Shuford 2008). The entire project site provides suitable habitat during migration for foraging, but there is no suitable nesting habitat on the project site. No Vaux's swifts were observed during surveys.

#### **4.1.18 Mountain Plover: SSC, BCC**

Mountain plover (*Charadrius montanus*) is found in semi-arid plains, grasslands, and plateaus. It uses open grasslands, plowed fields with little vegetation, and open sagebrush areas. Winter habitats include desert flats, and plowed fields. Mountain plovers are insectivores, feeding primarily on large ground-dwelling insects, including grasshoppers, beetles, and crickets (Shuford 2008). Their distribution was

modeled as occurring in the Chuckwalla Valley (CEC 2014a). The entire project site provides suitable habitat during migration but is unlikely to support suitable nesting habitat, since the Project site is outside its breeding range. No mountain plovers were observed on the Project site during surveys.

#### **4.1.19 Northern Harrier: SSC**

Northern harrier (*Circus cyaneus*) inhabits most of California at various times of the year, found up to 3000 m elevation. Northern harriers frequent meadows, grasslands, open rangelands, desert sinks, fresh and saltwater emergent wetlands. It is a widespread winter resident and migrant in suitable habitat. They primarily feed on small mammals, birds, frogs, small reptiles, crustaceans, and insects (Zeiner 1990). There is suitable foraging throughout the Project site, but no suitable nesting habitat. One northern harrier was observed in flight incidentally on the Project site during surveys, but a data point was not collected for the observation.

#### **4.1.20 Yellow-breasted Chat: SSC**

The yellow-breasted chat (*Icteria virens*) is an uncommon summer resident and migrant in coastal California, in foothills of the Sierra Nevada, and within the Colorado Desert. Breeding occurrences closest to the Project site are known from the Salton Sea and Colorado River. In southern California, yellow-breasted chats breed locally on the coast, and very locally inland (Garrett and Dunn 1981). During migration, they may be found in lower elevations of mountains in riparian habitat (McCaskie et al. 1979; Shuford 1990). The yellow-breasted chat may be found on the Project site during migration (most likely within desert dry wash woodland areas), but suitable nesting habitat is not present. No yellow-breasted chats were observed on the Project site during surveys.

#### **4.1.21 Crissal Thrasher: SSC**

Crissal thrasher (*Toxostoma crissale*) is a year-round resident of southeastern deserts, occupying dense shrubs in desert riparian and desert wash habitats, including mesquite, ironwood, and acacia. It primarily forages on the ground, feeding on invertebrates, berries, and seeds (Bent 1948; Shuford 2008). The Project site provides limited but suitable nesting and foraging habitat primarily associated with dry wash woodlands (see Figure 5). No crissal thrashers were observed within the Project during surveys.

#### **4.1.22 Elf Owl: BLMS, BCC**

Elf owl (*Micrathene whitneyi*) is found in lowland habitats that provide cover and good nesting cavities. It is most common farther east and north, in deserts with many tall saguaro cactus or large mesquites, and in canyons in the foothills, especially around sycamores or large oaks. The project site is near the western margin of its geographic range (Garret and Dunn 1981). The nearest nesting occurrence is near the Corn Springs campground and Cottonwood Springs vicinities (CNDDDB 2020). Elf owls are more common and widely distributed outside of California and probably have never been common in California due to limited geographic range and generally marginal habitat. The elf owl is migratory, spending winters in Mexico and southward. It arrives in California by March, and its breeding period extends from April to mid-July (Gould 1987).

The elf owl is a secondary cavity nester (it nests in cavities of trees and cacti, generally in disused woodpecker nests). Its nesting habitat is closely correlated with nesting habitat of woodpeckers, including Gila woodpecker (Hardy et al. 1999; Johnsgard 2002). Gila woodpeckers sometimes nest in



blue palo verde and palms, and elf owls have been documented nesting in blue palo verde near Wiley's Well, east of the project site, by Robert McKernan (former Director, San Bernardino County Museum; SBCM 2012a). Trees within the desert dry wash woodland habitat could provide suitable marginal habitat for nesting since desert dry wash woodland and larger trees are limited (see figure 5). No elf owls were observed on the Project site during surveys.

#### **4.1.23 Other listed Avian Species**

No suitable breeding or wintering habitat for the avian species below occur within or near the Project area. These state or federal listed bird species have been recorded at other utility-scale solar energy facilities. There is a moderate potential for them to pass within the Project vicinity during migration periods, but there is no suitable nesting or foraging habitat on the site for these species. None of these species were observed on the Project site during field surveys.

##### Yuma Ridgway's Rail: ST, CFP, FE

Yuma Ridgway's rail (*Rallus obsoletus yumanensis*), formerly known as Yuma clapper rail (*Rallus longirostris yumanensis*), nests in freshwater marshes. It is found along the lower Colorado River southward to its terminus at the Sea of Cortez, along the Gila River drainage in Arizona, at Lake Mead (and the Overton Arm) and its local tributaries, along the Virgin River in Nevada and Utah, and at the Salton Sea/Imperial Valley areas of California (CEC et. al 2014; USFWS 2014). Harrity and Conway (2019) captured 444 rails from 2016-2019 and attached transmitters to 103 rails to document annual migration and dispersal behaviors. As of December 16, 2019, they documented 24 migratory or dispersal movements (Harrity and Conway 2019). Yuma Ridgway's rail were thought to be mostly sedentary (Eddleman 1989), but recent rail mortalities at solar energy facilities and preliminary results of Harrity and Conway's (2019) telemetry study suggest that these rails fly over desert regions during dispersal and migration (Kagan et al. 2014, Harrity and Conway 2018). The transmitter data from this study confirms that rails migrate primarily at night (Harrity and Conway 2019). Most rails do not appear to follow the Colorado River corridor during migration, rather they cross vast expanses of desert upland and even open water to reach wintering grounds (Harrity and Conway 2019). These results help explain how Yuma Ridgway's rails perished at solar facilities far removed from any major sources of water or rail habitat (Kagan et al. 2014.) Outlier observations have been documented at Harper Dry Lake, East Cronese Dry Lake, and Desert Center, all at a great distance from known breeding areas (CNDDDB 2020).

##### Southwestern Willow Flycatcher: SE, FE

Southwestern willow flycatcher (*Empidonax traillii extimus*) breeds in dense riparian habitats in the southwestern United States, and winters in southern Mexico, Central America, and northern South America (USFWS 2002). The willow flycatcher species is comprised of several recognized subspecies, including the southwestern willow flycatcher, which is the only subspecies that nests in the region. The closest known breeding habitat to the Project site is approximately 35 miles away along the Colorado River and adjacent to the Salton Sea (CNDDDB 2020). Recent studies indicate that southwestern willow flycatchers do not migrate over the area of the desert where the Project site is located (BLM 2017). However, other willow flycatcher subspecies (not listed as threatened or endangered) may pass through

the area during migration. There is no suitable breeding habitat on the Project site and is outside the southwestern willow flycatcher's migratory routes.

Yellow billed cuckoo: SE, FT, BCC, BLMS

Western yellow-billed cuckoo (*Coccyzus americanus occidentalis*) breeds in expansive riparian areas in portions of California, Nevada, Arizona, and New Mexico. The closest known breeding habitat is located approximately 35 miles away along the Colorado River (CNDDDB 2018). During migration, western yellow-billed cuckoos migrate across the desert and use shrubland habitats, but there have been no documented sightings of western yellow-billed cuckoo within the Development Focus Areas (DFAs) identified in the DRECP LUPA (USFWS 2016). No suitable nesting habitat is present on the Project site, although it is possible that western yellow-billed cuckoo could occur on the site briefly during migration season.

Least Bell's Vireo: SE, FE

Least Bell's vireo (*Vireo bellii pusillis*) breeds in riparian habitats in southern California and portions of northern Baja California, Mexico and winters in southern Baja California, Mexico (USFWS 1998). Its numbers and distribution have probably increased since its listing, although it remains absent from large parts of its former range (USFWS 2016). The closest known breeding habitat to the site is to the northwest in the Big Morongo Canyon (USFWS 2020). Least Bell's vireos are also uncommon breeders at the Anza-Borrego Desert State Park, located approximately 70 miles southwest (USFWS 2016). The subspecies Arizona Bell's vireo (*V. b. arizonae*) is not ESA-listed, but is State-listed in California as endangered, and occurs along the lower Colorado River, approximately 35 miles east of the Project site.

Although there is little information on its migration behavior (USFWS 2016), least Bell's vireo likely migrates through the Colorado Desert. It is presumed that it may use riparian habitat and possibly upland scrub habitat during migration (USFWS 2016). No suitable nesting habitat is present on the Project site, although least Bell's vireo could occur on the site briefly, during migration season.

Avian Counts

A total of thirty avian species were observed when avian counts were conducted during spring surveys in the mornings. Appendix A-4 summarizes all avian counts observed during the survey period.

**4.1.23 American Badger: SSC**

The American badger is associated with dry open forest, shrub, and grassland communities with an adequate burrowing rodent population and friable soils. Badgers generally are associated with treeless regions, prairies, parklands, and cold desert areas (Zeiner et al. 1990). Badgers inhabit burrows and often prey on small mammals that inhabit burrows, as evidenced by claw marks along the edges of burrows. Suitable habitat exists for American badgers throughout the Project site. There are several canid burrows and complexes observed that could be used by the species although no badgers were observed and none of the burrows showed definitive badger sign.

#### **4.1.24 Desert Kit Fox: CPF**

Desert kit fox (*Vulpes macrotis arsipus*) is protected by the California Code of Regulations (Title 14, CCR: §460) and Fish and Game Code Section 4000 as a fur-bearing mammal. Title 14 of the California Code of Regulations, Section 460, stipulates that desert kit fox may not be taken at any time. Desert kit fox is a fossorial mammal that occurs in arid open areas, shrub grassland, and desert ecosystems within the Mojave and Sonoran Deserts. Desert kit fox typically occurs in association with its prey base, which includes small rodents, primarily kangaroo rats, rabbits, lizards, insects, and in some cases, immature desert tortoises (Zeiner et al. 1990). Burrow complexes that have multiple entrances provide shelter, escape, cover, and reproduction, but desert kit fox may utilize single burrows for temporary shelter. Litters of one to seven young are typically born in February through April (McGrew 1979).

During surveys, one old partial carcass of a kit fox was observed. Active and inactive desert kit fox burrows and scat were observed throughout the Project site (Figure 11, Appendix A-5). A total of twenty-five kit fox burrows were observed (nine of which showed active sign). These numbers will likely change over time since kit fox distribution is dynamic and change under natural conditions due to prey availability and other environmental factors such as the presence of coyotes that prey on kit fox pups.

#### **4.1.25 Desert Bighorn Sheep BLMS**

The desert bighorn sheep (*Ovis canadensis nelsoni*, also called Nelson's bighorn sheep) is found from the Peninsular and Transverse Ranges through most of the desert mountain ranges of California, Nevada, and northern Arizona to Utah. The Project site is well outside the range of the listed threatened Peninsular bighorn sheep, which was formerly recognized as a subspecies and now considered a distinct vertebrate population segment of the desert bighorn sheep. Essential habitat for desert bighorn sheep includes steep, rocky slopes of desert mountains, and areas where surface water is available during dry seasons. In the spring, when annual plants are available, bighorn sheep tend to disperse downhill to bajadas and alluvial fans to forage.

Habitat in the desert mountain ranges surrounding the upper Chuckwalla Valley is occupied by desert bighorn sheep, and they occasionally use the valley floor habitat either for foraging (near the lower mountain slopes) or as movement routes among mountain ranges. Due to the project's location on the valley floor near sites with comparable land uses and human activity patterns, the project is not likely to affect bighorn sheep behavior or habitat use to any large extent. No sign or evidence of desert bighorn sheep was found during field surveys, but scat is often difficult to distinguish from burro deer. Potential for occurrence is low.

#### **4.1.26 Burro Deer: CPGS**

Burro deer (*Odocoileus hemionus eremicus*) is a subspecies of mule deer (*Odocoileus hemionus*) that inhabits desert dry wash woodland communities in the Colorado region of the Sonoran Desert near the Colorado River. Some burro deer are year-around residents along the Colorado River, while others are transient and move between mesic and arid desert areas in response to seasonal water and forage availability. During hot summers burro deer concentrate along the Colorado River or the Coachella Canal where water developments have been installed and where microphyll woodland is dense and provides good forage and cover. With late summer thundershowers and cooler temperatures, burro deer move

away from the Colorado River and Coachella Canal into larger washes or wash complexes in the foothills and nearby mountains (BLM CDD 2002). Burro deer scat and tracks were observed throughout the Project site, but less so in areas with aeolian sand transport (Figure 11, Appendix A-5). Burro deer may be crossing the site to access nearby agriculture for water sources.

#### **4.1.27 Special Status Bats**

Bat roosts occur in the vicinity of the Project site in the McCoy Mountains approximately 20 miles east, Eagles Nest Mine within the Little Maria Mountains approximately 20 miles northeast, and Paymaster Mine within the Pinto Mountains approximately 30 miles (Larry LaPre, BLM, pers. comm.; CEC 2010). No active bat roosts were documented on the Project site during any of the surveys to date. It is not expected that any special status bat species would have a substantial roost on the Project site since habitat features most associated with these species (e.g. rock ledges, cliffs, large tree hollows, mine shafts) do not occur on the Project site. However, marginal roosting opportunities for bat species, such as the common canyon bat and California myotis, are available in tree cavities and soil crevices within dry desert wash woodland habitat, though limited. Additionally, suitable foraging habitat for common and special status bats is found on the Project site within desert dry wash woodland and near adjacent agricultural parcels where water may be available year-round.

Seven special status bat species may forage on or near the Project site; they are discussed further below. Suitable, but limited, roosting habitat may occur for several of these species within the dry wash woodland habitat and in nearby areas such as freeway underpasses. Those that occur closest to the Project vicinity, where camera stations were established (see section 3.2.7, figures 11, 12), did not show any sign of bat roosts at the time cameras were setup. Other special status bat species known from the region typically inhabit rocky sites and would not be expected to use the Project site for roosting. No bats were observed during surveys on the Project site.

##### Townsend's Big-Eared Bat: SSC, BLMS

Townsend's big-eared bat (*Corynorhinus townsendii*) roosts in caves, mines, abandoned dwellings, and large basal hollows of large trees (e.g., redwoods). Townsend's big-eared bat occurs from sea level to approximately 9,000 feet elevation within a range of habitats. It typically forages along streams and within woodlands. The Project site does not provide roosting areas for Townsend's big eared bat but does have foraging habitat in desert dry wash woodland and access to artificial water sources at agricultural farms nearby.

##### California Leaf-Nosed Bat: SSC, BLMS

California leaf-nosed bat (*Macrotus californicus*) occurs in the deserts of California, southern Nevada, Arizona and south to northwestern Mexico. In California, it is known from eastern San Bernardino, Riverside, and San Diego counties and all of Imperial County (CEC 2012). California leaf-nosed bat relies on caves and mines for roosting habitat. Foraging habitat typically consists of riparian and desert wash habitats, which occur on the Project. California leaf-nosed bat may forage within the Project site, but it is not expected to roost due to absence of suitable caves and mines.

#### Pallid Bat: SSC/BLMS

The pallid bat (*Antrozous pallidus*) is a locally common species throughout California, and a year-round resident in most of the range. It occupies a wide variety of habitats at elevations less than 6,000 feet including grasslands, shrublands, woodlands, and forests, and is most common in open, dry habitats with rocky areas for roosting; pallid bat roosts in cliffs, caves, crevices, mines, hollow trees, and various human-made structures (Zeiner 1990). The Project site may provide marginal suitable foraging habitat for pallid bats within the dry wash woodland but does not provide suitable roosting habitat. Acoustic bat surveys for Palen Solar Power Project (about 1 mile east of the Project site) detected pallid bat within the Project vicinity.

#### Western Mastiff Bat: SSC, BLMS

The western mastiff bat (*Eumops perotis californicus*) is widespread throughout the southwest U.S. and into Mexico. Its distribution in California is widespread, with year-round occurrence data primarily in central and southern California (Zeiner 1990). The western mastiff bat is found in a range of habitats, including coastal, forests, woodland, and desert scrub areas where roosting sites are available (Pierson and Rainey 1998). Roosting habitat typically consists of rocky crevices in canyons and cliffs with vertical or nearly vertical walls. The majority of roost sites are at least two meters above the ground (e.g., on cliff faces) and lacking obstructions. Suitable habitat for foraging occurs throughout the Project site, but roosting habitat is lacking. Western mastiff bat was detected within the vicinity on acoustic bat surveys for Palen Solar Power Project.

#### Western Yellow Bat: SSC

The western yellow bat (*Lasiurus xanthinus*) is a CDFW Species of Special Concern. It is found in Arizona, New Mexico, Mexico, and year-round in California. It is found in arid regions, in riparian, desert riparian, desert wash and palm oasis habitat. The western yellow bat is insectivorous, and roosts and feeds in palm oases and riparian habitats (Zeiner 1990). Potential roosting habitat, though marginal, exists within the Project site in areas where desert dry wash woodland exists. Suitable habitat for foraging also occurs in those same areas. Western yellow bat was detected within the vicinity during acoustic bat surveys for the Palen Solar Power Project.

#### Big Free-Tailed Bat: SSC

The big free-tailed bat (*Nyctinomops macrotis*) is distributed in the southwest U.S., and northern South America, generally from sea level to 8,000 feet in elevation. It is rare in California, prefers rocky terrain, and roosts in tree cavities and man-made structures. It wanders in autumn, out of its normal range (Zeiner 1990). Marginal foraging and potential roosting habitat for the big free-tailed bats occurs within the Project in desert dry wash woodland. Big free-tailed bat was detected within the Project vicinity through acoustic surveys conducted for Palen Solar Energy Project.

#### Pocketed Free-Tailed Bat: SSC

The pocketed free-tailed bat (*Nyctinomops femorosaccus*) is common in Mexico but less common in western North America, from southern California, central Arizona, southern New Mexico, and western

Texas (WBWG 2018). The pocketed free-tailed bat has been documented in Riverside, San Diego, and Imperial counties. Typical habitats include pinyon-juniper woodlands, desert scrub, desert succulent shrub, desert riparian, desert wash, alkali desert scrub, Joshua tree, and palm oasis. Roosting habitat typically includes rock crevices associated with granite boulders, cliffs, or rocky canyons at a height suitable for approach and takeoff (CNDDDB 2020). Pocketed free-tailed bats occur in the desert from March through August, when they then migrate out of the area (BLM 2011). Suitable habitat for foraging exists on the Project site, but roosting habitat is lacking. Call sequences that may have been pocketed free-tailed bat were detected within the Project vicinity during acoustic surveys for Palen Solar Energy Project but lacked features for definitive confirmation.

## **4.2 Wildlife Movement**

Wildlife movement among habitat areas is a part of regular activities and may be needed for long-term population sustainability. Land use changes can impact wildlife movement across the landscape, leading to habitat fragmentation and population isolation. When habitat is converted to other uses, it separates or isolates the remaining habitat areas, which creates less habitat availability, and less opportunity for wildlife to make use of the remaining habitat, due to its physical isolation. Habitat areas may be isolated from one another by distance across unfavorable habitat, or by linear barriers such as roadways or aqueducts. Barriers may be impassable for some species, such as a wide busy road, for a slow-moving animal, or may be only minor interruptions to movement, such as a narrow, lightly travelled road. Fragmentation and subsequent population isolation can affect wildlife populations by limiting dispersal and genetic exchange, limiting movement within the home-ranges for wide-ranging species, and limiting the opportunity for populations to occupy new habitat in response to the effects of climate change. Fragmentation also increases habitat “edge” (i.e., habitat adjacent to other land uses), leading to increased exposure to invasive species, human disturbance (vehicles, trash dumping, etc.), and an overall reduction of biodiversity and alteration or degradation of ecological processes. Within the Project vicinity are existing and developing solar farms, other commercial developments, and agriculture that contributes to the fragmentation of habitat (Figure 12).

Accessibility between habitat areas (i.e., “connectivity”) is important to long-term genetic diversity and demography of wildlife populations. In the short term, connectivity may also be important to individual animals’ ability to occupy their home ranges, if their ranges extend across a potential movement barrier. These considerations apply to greater or lesser extent to all plants and animals. Plant populations “move” over the course of generations via pollen and seed dispersal; most birds and insects travel and disperse via flight; terrestrial species, including small mammals, reptiles, arid land amphibians, and non-flying invertebrates, disperse across land. Therefore, landscape barriers and impediments are more important considerations for movement of terrestrial species. These considerations are especially important for rare species and wide-ranging mammals, which both tend to exist in lower population densities.

In developed landscapes where remnant habitat exists as partially isolated patches surrounded by other land uses, planning for wildlife movement generally focuses on “wildlife corridors” to provide animals with access routes between habitat patches. In largely undeveloped areas, including the Chuckwalla Valley, wildlife habitat is available in extensive open space areas throughout much of the region, but

specific barriers may impede or prevent movement. In these landscapes, wildlife movement planning focuses on specific sites where animals can cross linear barriers (e.g., wash crossings beneath Interstate 10), and on broader linkage areas that may support stable, long-term populations of target species and allow demographic movement and genetic exchange among populations in distant habitats (e.g., surrounding mountains).

The California Desert Connectivity Project provides a comprehensive and detailed habitat connectivity analysis for the California deserts (Penrod et al. 2012). The Connectivity Project identified a Desert Linkage Network to maintain habitat for movement between landscape blocks. The landscape blocks identified in the project vicinity are the Palen–McCoy Mountains to the northeast and the Chocolate Mountains to the southwest. Broad habitat linkages connect these landscape blocks. The DRECP identifies a wide multi-species linkage area southwest of the Project site, within which a narrower wildlife corridor will be defined (Figures 1 and 12). The Project site is not located within any multi-species linkage area. The final design of the project will follow all CMA requirements and may have a reduced footprint that will avoid or have a reduced footprint within multi-species linkage boundaries.

In the Chuckwalla Valley, the biologically important functions of large mammal movement are the long-term demographic and genetic effects of occasional animal movement among mountain ranges and other large habitat areas. Animals such as desert bighorn sheep may travel across the valley infrequently, to reach other subpopulations in surrounding mountains. In contrast to large animal movement, desert tortoises and other less-mobile animals may live out their entire lives within a linkage area between larger habitat blocks; for these species, movement among surrounding habitat areas may take place over the course of several generations.

Movement opportunity varies for each species, depending on motility and behavioral constraints, as well as landscape impediments. For many terrestrial wildlife species, movement across the Chuckwalla Valley, including movement to and from the project site, or across the site, is limited by anthropogenic barriers or land uses. The I-10 freeway, located south of the project site, is a significant obstruction to movement by terrestrial wildlife. Some species, such as coyote, may learn to cross the freeway safely. However, the freeway presents an impassable or high-risk barrier to north-south movement for most terrestrial species. Other linear features, such as smaller paved and unpaved roads and transmission lines have only minimal effects on wildlife movement.

On the 32-mile stretch of I-10 between the Desert Center and Wiley Wells Road exits there are 24 crossings that provide safe access under the freeway (CEC, 2010). Other than these crossings, the freeway is a nearly complete barrier to north-south terrestrial wildlife movement in the Chuckwalla Valley. A survey of potential tortoise accessibility across the I-10 investigated these 24 crossings (oriented approximately in a north-south direction) for suitability for large mammals, small mammals, and reptiles (CEC, 2010). The survey found that fencing was often missing or in disrepair, was not tethered to the underpasses, and does not function to funnel wildlife under the interstate. The study concluded the underpasses provide connectivity and safe movement corridors between habitat areas to the north and south of the I-10, but the fencing does not prevent animals from accessing I-10. Wildlife species and sign detected at the undercrossings included lizards, rodents, rabbit, roadrunner, ground squirrel, fox, coyote, bobcat, and burro deer. Additionally, the CDFW and USFWS have documented

burro deer using I-10 undercrossings. Six of these undercrossings are located within the multi-species linkage area (Figure 12).

Wildlife cameras set up for the duration of one month during spring 2020 season at seven of these I-10 undercrossings near the Project site, identified several species using those corridors. At least five of these camera stations were within the multi-species linkage area. Table 6 below summarizes the species detected by the wildlife cameras using these undercrossings (Figure 12, Photos 7-10).

**Table 6.** Summary of Wildlife Camera Observations

Species Observed	Camera Station Number						
	C08	C09	C10	C11	C12	C13	C14
burro deer ( <i>Odocoileus hemionus eremicus</i> )			x	x	x	x	
bobcat ( <i>Lynx rufus</i> )					x		
coyote ( <i>Canis latrans</i> )	x		x		x	x	x
desert kit fox ( <i>Vulpes macrotis</i> )	x						x
black-tailed jackrabbit ( <i>Lepus californicus</i> )			x		x	x	x
desert wood rat ( <i>Neotoma lepida</i> )			x	x			
round-tailed ground squirrel ( <i>Xerospermophilus tereticaudus</i> )							x
desert kangaroo rat ( <i>Dipodomys deserti</i> )					x		
rodent (unknown)			x	x	x		
avian species (unknown)			x	x	x		
hummingbird species ( <i>Trochilide sp</i> )			x				
desert iguana ( <i>Dipsosaurus dorsalis</i> )			x				
desert spiny lizard ( <i>Sceloporus magister</i> )				x			
western whiptail lizard ( <i>Aspidoscelis tigris</i> )				x			
Eurasian collared dove ( <i>Streptopelia decaocto</i> )						x	
insect ( <i>Lepidoptera sp</i> )		x			x		

### 4.3 Special Status Plant Species

Forty-two special status plant species were reviewed for their potential to occur within the Project site and its vicinity based on regional plans and database records (Appendix C). The probability of occurrence is defined as follows:

- Present: Species was observed at the time of the survey



- High: Both a historical record exists of the species within the project site or its immediate vicinity (approximately 5 miles) and the habitat requirements associated with the species occur within the project site.
- Moderate: Either a historical record exists of the species within the immediate vicinity of the project site (approximately 5 miles) or the habitat requirements associated with the species occur within the project site.
- Low: No records exist of the species occurring within the project site or its immediate vicinity and/or habitats needed to support the species are of poor quality.
- Minimal: Species was not observed during focused surveys conducted at an appropriate time for identification of the species, or species is restricted to habitats that do not occur within the project site

Special status species detected within the Project site or having moderate potential to occur based on the presence of suitable habitat are discussed further in this section. Special status species observed are identified in Appendix A-6 and mapped on Figure 13.

#### **4.3.1 Crucifixion Thorn: CRPR 2B.2**

Crucifixion thorn (*Castela emoryi*) has 177 records occurring within California. In Riverside County, several records are near or within Desert Center, including Desert Sunlight Solar Farm just north of the Project and at Athos Solar Project (CCH 2020). There is suitable habitat for crucifixion thorn within wash areas of the Project site. It was observed at two locations at the Project site (Figure 13). It is a large conspicuous shrub and can be located and identified at any time of year, even in a year of poor rainfall.

#### **4.3.2 Glandular Ditaxis: CRPR 2B.2**

Glandular ditaxis (*Ditaxis claryana*) is an annual or short-lived perennial that blooms in the fall following the start of the rainy season that occurs in Sonoran desert scrub. There are 49 occurrences in the Consortium of California Herbaria (CCH 2020) and there is one record within Desert Center and another near Corn Springs ACEC, south of I-10 (CNDDDB 2020). Suitable habitat occurs within the Project site, but it was not observed during plant surveys.

#### **4.3.3 California Ditaxis: CRPR 3.2**

California ditaxis (*Ditaxis serrata* var. *californica*) has a CRPR of 3.2 and a NatureServe rank of G3G4/S2 S, which indicates more information is needed about the status of this species. California ditaxis may be a glabrous variety of the common *Ditaxis neomexicana* (CEC 2010). It occupies Sonoran Desert scrub vegetation and prefers sandy washes and alluvial fans of the foothills and lower desert slopes, from 100 to 3,000 feet amsl. It is known to occur in San Bernardino, Riverside, Imperial, and San Diego counties of California and in Sonora, Mexico (CNPS 2020). There are 45 records of this species in California, primarily from Riverside County (CCH 2020). There is suitable habitat on the Project site, but it was not observed during surveys.

#### **4.3.4 Utah Milkvine: CRPR 4.2**

Utah milkvine (*Cynanchum utahense* [= *Funastrum utahense*]) has 149 records from the Consortium of California Herbaria database primarily from San Bernardino and San Diego counties, but there are also several records in Riverside County. There is one record of this species north of Desert Center and another record just southwest of Palen Lake. This twining perennial occurs in sandy, gravelly Mojavean

desert scrub. Suitable habitat exists throughout the Project site, but Utah milkvetch was not observed during surveys.

#### **4.3.5 Desert Unicorn Plant: CRPR 4.3**

Desert unicorn plant (*Proboscidea althaeifolia*) has limited distribution but is not very threatened in California. It is a low-growing, perennial species that occurs in sandy washes within Sonoran desert scrub vegetation in San Bernardino, Imperial, Riverside, and San Diego counties of California. There are 36 records in Riverside County, several of which are from the Chuckwalla Mountains and Desert Center area (CCH 2020). It is a late-season bloomer (May to August) but has large and distinctive seed pods that can be detected during the spring season and fleshy root structure that can remain dormant in dry years (BLM 2011). Suitable habitat occurs within the Project site, but it was not observed during surveys.

#### **4.3.6 Chaparral Sand Verbena: BLMS, CRPR 1B.1**

Chaparral sand verbena (*Abronia villosa* var. *aurita*) has 238 records within several counties in southern California, many of which are in Riverside County. Its distribution and identification are unclear in published reference works, including Spellenberg (2002), CNPS (2020), and CNDDDB (CDFW 2020). It was added to the CNPS Inventory based on recommendations by Andrew C. Sanders of the UC Riverside Herbarium. The primary conservation concern is for chaparral sand-verbena occurrences in western Riverside County and other locations outside the desert (see Roberts et al. 2004). These western plants appear to be distinct from the very common desert sand verbena, *Abronia villosa* var. *villosa*. Plants in the low desert often match the characteristics of the western Riverside County populations, but they are not regionally rare. There is one record that is very close to the Project site, on the Palen sand dunes in the vicinity of the Desert Lily Sanctuary, located in 2012. Suitable habitat occurs within the Project site. No chaparral sand verbena were observed on the Project site during surveys. Only the more common hairy sand verbena (*Abronia villosa* var. *villosa*) was observed.

#### **4.3.7 Harwood's Milkvetch: CRPR 2B.2**

Harwood's milkvetch (*Astragalus insularis* var. *harwoodii*) has historic and recent collections that include Ogilby Road in Imperial County and three locales west of Blythe, the Pinto Basin, and Chuckwalla Basin in Riverside County. Harwood's milkvetch has also been reported from Baja California, Sonora Mexico, and portions of Yuma County, Arizona. Its primary habitat is windblown sand. There are several CNDDDB records for this species within the Project vicinity (CNDDDB 2020). Many new occurrences were documented in Chuckwalla Valley and the Palo Verde mesa during surveys for the Blythe Solar Power Project, the Genesis Solar Energy Project, McCoy Solar Energy Project, and Palen Solar Power Project study areas. There is suitable habitat for Harwood's milkvetch and two locations (four individuals) were observed on the northern portion of the Project site (Figure 13, Appendix A-6).

#### **4.3.8 Abram's Spurge: CRPR 2B.2**

Abram's spurge (*Chamaesyce abramsiana* [= *Euphorbia abramsiana*]) occurs in saline scrub flats, playas, and along inlets and floodplains of playas. There are 137 records in California within Imperial, Riverside, San Bernardino, and San Diego counties. The records within Chuckwalla Valley closest to the Project site were near Palen Dry Lake and Pinto Basin. Marginal suitable habitat is present within the saltbush scrub of the Project site, but none were observed during surveys.

#### **4.3.9 Ribbed Cryptantha: CRPR 4.3**

Ribbed cryptantha (*Cryptantha costata*) has 279 records from several locations throughout Riverside, Imperial, San Diego, and Imperial counties (CCH 2020). It occurs in windblown sand habitats. A large local population of ribbed cryptantha was observed just east of the proposed Palen Solar Power Project. Suitable habitat occurs at the Project site and several locations were identified that had large ribbed cryptantha populations clustered around those areas (Figure 13, Appendix A-6).

#### **4.3.10 Harwood's Eriastrum: CRPR 1B.2, BLMS**

Harwood's eriastrum (*Eriastrum harwoodii*), also commonly known as Harwood's woollystar, has a CRPR of 1B.2, has a NatureServe rank of G2/S2 and is a BLM sensitive species. It is a spring annual, typically found in dunes associated with the margins around dry lakes such as Dale, Cadiz, and Soda lakes (CNPS 2020). Reports of this species are known from San Bernardino, Riverside, Imperial, and San Diego counties in California and in Sonora, Mexico (CNPS 2020). There are 118 records of this species in California (CCH 2020). It has been observed within partially stabilized dunes at nearby project sites. Suitable habitat is present on the Project site and Harwood's eriastrum was observed at five locations within the Project site with several individuals at each location (Figure 13, Appendix A-6).

#### **4.3.11 Jackass Clover: CRPR 2B.2**

Jackass clover (*Wislizenia refracta* ssp. *refracta*) is commonly associated with sandy washes, roadsides, or alkaline flats. There are 28 occurrences in the Consortium of California Herbaria most of which are located in San Bernardino County near Twentynine Palms, with only one record in Riverside County east of Indio (CCH 2020). Jackass clover was also documented at several locations from the northern to southern end of Palen Lake in dune habitats during a detailed vegetation mapping and classification project conducted by CNPS Vegetation Program for BLM (Evens & Hartman 2007). Jackass clover is found in sandy washes, roadsides, or alkaline flats. Suitable habitat is present on the Project site, particularly in the northern and eastern boundaries, but it was not observed during surveys.

#### **4.3.12 Palmer's Jackass Clover: CRPR 2B.2**

Palmer's jackass clover (*Wislizenia refracta* ssp. *palmeri*) has 15 occurrences in the Consortium of California Herbaria with at least three records near Desert Center (CCH 2020). It typically occupies sandy washes, within Sonoran desert scrub vegetation. Suitable habitat is present throughout the Project site, but it was not observed on the Project site during surveys.

#### **4.3.13 Creosote Bush Rings: BLMS**

No creosote bush rings were observed on the Project site during surveys.

#### **4.3.14 Cacti, Yucca, and Native Trees**

Native cacti, succulents, and trees are generally not ranked as special status plant species but the harvesting of these native plants is regulated under the California Native Plant Protection Act (Fish and Game Code §§ 1900-1913) and the California Desert Native Plant Act of 1981 (Food and Agricultural Code § 80001 et. seq.; Fish & Game Code §§ 1925-1926). Any vegetation to be salvaged and removed from the site (such as cactus or yucca) would be subject to sale at appraised value, according to CFR

43:5420.0-6. If the cacti or yucca is salvaged and/or transplanted offsite, as approved by BLM, then this resource is not subject to sale but remains in BLM ownership. These species are:

- Fishhook cactus (*Mammillaria tetrancistra*)
- Ocotillo (*Fouquieria splendens* ssp. *splendens*)

Additionally, five species of native trees were found within the Project site:

- Desert ironwood (*Olneya tesota*)
- Blue palo verde (*Parkinsonia florida*)
- Honey mesquite (*Prosopis glandulosa*)
- Smoke tree (*Psoralea arguta*)
- Catclaw acacia (*Senegalia greggii*)

#### **4.4 Invasive Weeds**

Invasive weeds are non-native (exotic) plants included on the weed lists of the California Invasive Plant Council (Cal-IPC), or those weeds of special concern identified by the BLM. There are also some weeds designated as “noxious” by California Department of Food and Agriculture (CDFA) or the U.S. Department of Agriculture. Invasive weeds are of concern in wild lands because of their potential to degrade habitat and disrupt the ecological functions (Cal-IPC 2020). The following invasive weeds were identified on the Project site during Ironwood’s field surveys and summarized in Appendix A-7(Figure 13).

##### Sahara Mustard (*Brassica tournefortii*)

Sahara mustard has a highly invasive rating on Cal-IPC (Cal-IPC 2020). It has severe ecological impacts on physical processes, plant and animal communities, and vegetation structure, as well as having reproductive biology and other attributes that are conducive to moderate to high rates of dispersal and establishment (Cal-IPC 2020). Sahara mustard is native to the deserts of North Africa, the Middle East, and the Mediterranean regions of southern Europe (Bossard et al. 2000). Initial establishment of this species in California occurred through the importation of date palms from the Middle East to the Coachella Valley during the early 1900s (Bossard et al. 2000). Sahara mustard currently occurs across Riverside County, as well as all neighboring counties (Cal-IPC 2020). Sahara mustard was found throughout the Project site during surveys

##### Russian Thistle (*Salsola tragus*)

Russian thistle has a Limited-to-Moderate rating by the Cal-IPC, indicating a species that is invasive but has an ecological impact that is minor on a statewide level, or there was not enough information to justify a higher score. Its reproductive biology and other attributes result in low to moderate rates of invasiveness. Ecological amplitude and distribution are generally limited, but it may be locally persistent and problematic. Russian thistle is listed on the CDFA Noxious Weed List, making it subject to state laws and regulations regarding its spread and pollution of an area (CDFA 2020). Russian thistle is an annual herb that is found in open and disturbed areas in the Mojave Desert and throughout western North America (MacKay 2003). Otherwise known as tumbleweed, it becomes large and round with age, the

dried plant breaks off and rolls with the wind to aid in seed dispersal. Native to Eurasia, this plant was likely introduced around the turn of the century. It typically occurs on sandy soils on disturbed sites, cultivated and abandoned fields, and disturbed natural and semi-natural plant communities (CDFA 2020). Russian thistle was observed on the edges of the Project site that were closer to adjacent fallow agriculture.

#### Redstem filaree (*Erodium cicutarium*)

Redstem filaree has a limited invasive rating on Cal-IPC (Cal-IPC 2018) and is not listed on the CDFA Noxious Weed List (CDFA 2020). This species is an aggressive annual/biannual of the family Geraniaceae (geranium) family that is very widespread throughout California and is commonly found along roadsides, grasslands, fields, and semi-desert areas. It occurs throughout the Project site and often carpets large areas, out-competing native grasses and forbs. It occurs throughout the Project site.

#### Mediterranean grass (*Schismus barbatus* and *S. arabicus*)

Mediterranean grass has a limited invasive potential (CAL-IPC 2020) and is not listed by CDFA. It is an annual grass found in both central and southern California, particularly in disturbed areas and deserts, probably introduced at the turn of the century (CDFA 2020). It contributes to increased fire ignition and spread due to accumulation of dry thatch during dry seasons. Wildfire, in turn, contributes to the type-conversion of desert shrubland into annual grassland. These species' reproductive biology and other attributes result in low to moderate rates of invasiveness. Spread may occur from seed dispersal associated with soil disturbance, vegetation cutting, and from vehicle tires and footwear. Increase of these species is most likely to occur in areas where it already exists. Mediterranean grass is prevalent throughout Sonoran creosote bush scrub of the Project site. BLM and other agencies recognize that because of its widespread distribution, Mediterranean grass is not feasible to eradicate. It occurs throughout the Project site.

#### London rocket (*Sisymbrium irio*)

London rocket has a moderate rating by the Cal-IPC, indicating that the species has substantial and apparent, but generally not severe, ecological impacts on physical processes, plant and animal communities, and vegetation structure. Ecological amplitude and distribution may range from limited to widespread. It is a winter annual forb/herb (family Brassicaceae), which can be found in abandoned fields, waste places, roadsides, and orchards. It matures earlier in the year than native species, allowing it to out-compete them. It is not listed on the CDFA noxious weed list. The species distribution is generally spreading in California desert regions, (Cal-IPC, 2020). During surveys, this species was generally limited to areas directly underneath desert ironwood (*Olneya tesota*) trees or other large shrubs on the edges of the Project site.

#### Cheatgrass (*Bromus tectorum*)

Cheatgrass has an invasive rating of high according to Cal-IPC and is not listed on the CDFA noxious weed list. It has severe ecological impacts on physical processes, plant and animal communities, and vegetation structure. Its reproductive biology and other attributes are conducive to moderate to high

rates of dispersal and establishment. Cheatgrass is an annual grass (family Poaceae). It is the dominant grass on sagebrush (*Artemisia* species) rangelands on the Modoc Plateau in northeastern California and along the eastern Sierra Nevada to Owens Valley. This weed overcrowds native grasslands and croplands. Replacement of native grasses by cheatgrass increases the frequency and extent of wildfires. Cheatgrass has documented low to moderate abundance across much of the California deserts (Cal-IPC, 2020). This grass was observed in low abundance on the edges of the Project site.

#### Non-native, naturalized species

Other non-native plant species observed on the Project that are not considered invasive but have become naturalized include:

- Sowthistle (*Sonchus oleraceus*)
- Spiny sowthistle (*Sonchus asper*)
- Hedge mustard (*Sisymbrium orientale*)

## 5.0 REFERENCES

- AECOM. 2009. Palen Solar Power Project Biological Technical Report. Riverside County, California. Submitted to Solar Millennium, LLC, Berkeley, California, and Chevron Energy.
- AECOM. 2010. Fall Botanical Surveys. Palen Solar Power Project. CEC Docket No. 09-AFC-7. TN 58879. May 17, 2010.
- Andersen, M. C., J. M. Watts, J. E. Freilich, S. R. Yool, G. I. Wakefield, J. F. McCauley and P. B. Fahnestock. 2000. Regression-tree modeling of desert tortoise habitat in the central Mojave Desert. *Ecological Applications* 10(3): 890-900.
- Andre, Silverman, pers. comm. 2010. Cited in CEC Revised Staff Assessment, Palen Solar Project, Part 2.
- Aspen Environmental Group (Aspen). 2019. Project Memorandum – Arica Solar Project and Victory Pass Solar Project: Biological Resources Survey Work Plan. September 23, 2019.
- Averill-Murray, Roy C., C. Darst, N. Strout, and M. Wong. 2013. Conserving Population Linkages for The Mojave Desert Tortoise (*Gopherus agassizii*). *Herpetological Conservation and Biology* 8(1):1-15. Published: 30 April 2013.
- Baldwin, B.G., D.H. Goldman, D.J. Keil, R. Patterson, T.J. Rosatti, and D.H. Wilken, editors. 2012. The Jepson manual: vascular plants of California, second edition. University of California Press, Berkeley.
- Barrett, S. L. 1990. Home range and habitat of the desert tortoise (*Xerobates agassizii*) in the Picacho mountains of Arizona. *Herpetologica* 46(2): 202-206.
- Beal, F. E. L. 1907. Birds of California in relation to the fruit industry, part 1. U.S. Dept. Agri. Biol. Surv. Bull. 30.
- Bent, A. C. 1948. Life histories of North American nuthatches, wrens, thrashers, and their allies. U.S. Natl. Mus. Bull. 195. 475pp.
- Berry, K.H. 1986. Desert tortoise (*Gopherus agassizii*) relocation: Implications of social behavior and movements. *Herpetological* 42:113-125.
- Bitter, Dave. 2010. Investigator's Annual Report, Joshua Tree National Park. Golden Eagle Survey Findings. JOTR-00157.
- Bitter, Dave. 2011. Investigator's Annual Report, Joshua Tree National Park. Golden Eagle Survey Findings. JOTR-00157.
- Bleich, V. C., J. D. Wehausen, R. R. Ramey II, and J. L. Rechel. 1996. Metapopulation theory and mountain sheep: implications for conservation. Pages 353-373 in D. R. McCullough, editor. *Metapopulations and wildlife conservation*. Island Press, Washington D.C., USA.
- Bloom Biological, Inc. (BBI). 2013a. (TN 200010) Palen Solar Electric Generating Facility Spring 2013 Avian Survey Results. Prepared for Palen Solar Holdings, Inc., Oakland, California. Prepared by BBI, Lake Forest, California. July 2013.
- Bloom Biological, Inc. (BBI). 2013b. Palen Solar Electric Generating Facility Summer 2013 Avian Survey Results. Prepared for Palen Solar Holdings, Inc., Oakland, California. Prepared by BBI, Lake Forest, California. August 2013.
- Bloom Biological, Inc. (BBI). 2013c. Palen Solar Electric Generating System 2013 Golden Eagle Nesting Survey Results. Prepared for Palen Solar Holdings, Inc., Oakland, California. Prepared by BBI, Lake Forest, California. October 2013.

- Bloom Biological, Inc. (BBI). 2013d. Palen Solar Electric Generating System Winter 2013 Golden Eagle Survey Results. Prepared for Palen Solar Holdings, Inc., Oakland, California. Prepared by BBI, Lake Forest, California. March 2013.
- Bloom Biological, Inc. (BBI). 2013e. (TN 70242) PSEGS Winter 2013 Golden Eagle Survey Results. Prepared for BrightSource Energy, Inc. March 2013.
- Bureau of Land Management California Desert District (BLM CDD) and California Department of Fish and Game Inland, Desert, and Eastern Sierra Region. 2002. Proposed Northern and Eastern Colorado Desert Coordinated Management Plan and Final EIS, July 2002.
- Bureau of Land Management (BLM). 2009. Survey Protocols Required for NEPA/ESA Compliance for BLM Special Status Plants.
- Bureau of Land Management (BLM). 2010. Special Status Animals in California, Including BLM Designated Sensitive Species.
- Bureau of Land Management (BLM). 2011. Plan Amendment and Final EIS for the Palen Solar Power Project. Palm Springs -South Coast Field Office. May 2011.
- Bureau of Land Management (BLM). 2012. Results of 2012 Golden Eagle Nesting Surveys of the California Desert and Northern California Districts. Prepared by BioResource Consultants, November 2012.
- Bureau of Land Management (BLM). 2013. Golden Eagle Nest Occurrences, DRECP. Dataset in Databasin.org . <https://databasin.org/datasets/9e7e4a80961f468887322ddd8e840777>. Last accessed July 2020.
- Bureau of Land Management (BLM). 2013. Draft Supplemental EIS for the Palen Solar Electric Generating System. Palm Springs -South Coast Field Office. July 2013.
- Bureau of Land Management (BLM). 2016. Desert Renewable Energy Conservation Plan: Land Use Plan Amendment and Final Environmental Impact Statement. Prepared by U.S. Bureau of Land Management in partnership with U.S. Fish and Wildlife Service, California Energy Commission, and California Department of Fish and Wildlife. BLM/CA/PL-2016/03+1793+8321. October 2015.
- Bossard, C.C. J.M Randall and M.C. Hoshovsky, 2000. Invasive Plants of California's Wildlands. University of California Press, Berkeley, CA.
- Brown, P. E., and W. E. Rainey. 2014. Bat Habitat Assessment for Palen Solar Electric Generation System. Conducted for Alice E. Karl and Associates. Conducted by Brown-Berry Biological Consulting. May 17, 2014.
- California Burrowing Owl Consortium (CBOC). 1993. Burrowing Owl Survey Protocol and Mitigation Guidelines. Prepared by the California Burrowing Owl Consortium. April 1993.
- California Invasive Plant Council (Cal-IPC). 2020. California Invasive Plant Inventory Database. Available at: <<http://www.cal-ipc.org/paf>> (Accessed July 2020).
- California Department of Food and Agriculture (CDFA). 2020. Amend Section 4500. Noxious Weed Species. [http://www.cdfa.ca.gov/plant/ipc/encycloweedia/encycloweedia\\_hp.htm](http://www.cdfa.ca.gov/plant/ipc/encycloweedia/encycloweedia_hp.htm). (Accessed July 2020).
- California Department of Fish and Game (CDFG). 2000. Guidelines for Assessing the Effects of Proposed Projects on Rare, Threatened and Endangered Plants and Natural Communities.
- California Department of Fish and Wildlife. (CDFW). 2012. Staff Report on Burrowing Owl Mitigation. March 7.
- California Department of Fish and Wildlife. (CDFW). 2020a. Natural Diversity Database. July 2020. Special Animals List. Periodic publication. 51 pp.



- California Department of Fish and Wildlife. (CDFW). 2020b. A Status Review of Townsend's Big-Eared Bat (*Corynorhinus townsendii*) in California.
- California Department of Fish and Wildlife. (CDFW). 2020c. Natural Diversity Database Special Vascular Plants, Bryophytes, and Lichens List. Quarterly Publication. 73 pp. July 2020.
- California Department of Fish and Wildlife. (CDFW). 2020d. California Sensitive Natural Communities. <https://www.wildlife.ca.gov/Data/VegCAMP/Natural-Communities>. July 2020.
- California Energy Commission (CEC). 2010. RSA (Revised Staff Assessment). Palen Solar Project, Part 2. September 2010. (TN 58497)
- California Energy Commission (CEC), California Energy Commission, California Department of Fish and Wildlife, U.S. Bureau of Land Management, U.S. Fish and Wildlife Service. 2014a. Draft Desert Renewable Energy Conservation Plan (DRECP) and Environmental Impact Report/Environmental Impact Statement. July 2020 [www.drecp.org](http://www.drecp.org).
- California Energy Commission (CEC), California Energy Commission, California Department of Fish and Wildlife, U.S. Bureau of Land Management, U.S. Fish and Wildlife Service. 2014b. Draft Desert Renewable Energy Conservation Plan (DRECP) and Environmental Impact Report/Environmental Impact Statement. Appendix Q, Baseline Biological Report.
- California Energy Commission (CEC), California Energy Commission, California Department of Fish and Wildlife, U.S. Bureau of Land Management, U.S. Fish and Wildlife Service. 2014c. Draft Desert Renewable Energy Conservation Plan (DRECP) and Environmental Impact Report/Environmental Impact Statement. Appendix B, Species Profiles.
- California Energy Commission (CEC), California Energy Commission, California Department of Fish and Wildlife, U.S. Bureau of Land Management, U.S. Fish and Wildlife Service. 2014d. Draft Desert Renewable Energy Conservation Plan (DRECP) and Environmental Impact Report/Environmental Impact Statement. Volume III Section 7, Environmental Setting/Affected Environment, Biological Resources.
- California Geological Survey. 2015. Comments and Additional Information for Incorporation into the Draft Desert Renewable Energy Conservation Plan (DRECP) and Environmental Impact Report/Environmental Impact Statement (EIR/EIS). February 5, 2015.
- California Native Plant Society (CNPS). 2007. Vegetation Survey and Classification for the Northern & Eastern Colorado Desert Coordinated Management Plan (NECO).
- California Native Plant Society (CNPS). 2020. CNPS Inventory of Rare and Endangered Plants, Riverside County. Available at: < <http://www.rareplants.cnps.org/> > Accessed July 2020
- California Natural Diversity Data Base (CNDDB). 2020. Aztec Mines, Corn Spring, East of Aztec Mines, East of Victory Pass, Ford Dry Lake, Palen Lake, Palen Mountains, Pilot Mountain, and Sidewinder Well 7.5 minute USGS quadrangles.
- Consortium of California Herbaria (CCH). 2020. Search results for taxa from CNPS Inventory in Riverside County, retrieved July 2020. Available at: <http://ucjeps.berkeley.edu/consortium/>.
- Corvus Ecological. Personal communication regarding Chuckwalla raptor surveys. August 2020
- Davis, F. and Soong, O. 2013. Mojave fringe-toed lizard - Species Distribution Model, DRECP. Bren School of Environmental Science & Management. University of California, Santa Barbara Conservation Biology Institute. Jun 25, 2013 (modified Dec 3, 2013).
- D'Antonio, C.M. 1990a. Invasion of coastal plant communities in California by the introduced iceplant, *Carpobrotus edulis* (Aizoaceae). Ph.D dissertation, University of California, Santa Barbara, CA .

- Desert Research Institute (DRI) 2013. Geomorphic Assessment of Sand Transport for the Modified Project (Palen Solar Electric Generating System). Draft Final Report. Prepared for Aspen Environmental Group. Prepared by Dr. N. Lancaster and Dr. T. Bullard, Division of Earth and Ecosystem Sciences and Dr. J. Gillies Division of Atmospheric Sciences. July 23, 2013.
- Duda, J. J., A. J. Krzysik and J. E. Freilich (1999). "Effects of drought on desert tortoise movement and activity." *Journal of Wildlife Management* 63(4): 1181-1192.
- Dunn, J. L., and Garrett, K. L. 1997. *A Field Guide to Warblers of North America*. Houghton Mifflin, Boston.
- EDAW AECOM. 2009. Palen Solar Power Project Burrowing Owl Technical Report. July 2009.
- EDAW AECOM and Bloom Biological, Inc. (BBI). 2009. Palen Solar Power Project Avian Point Count Survey Technical Report. Prepared for Solar Millennium, LLC and Chevron Energy Solutions. August 2009.
- Edwards, H.H., and G.D. Schnell. 2000. Gila Woodpecker (*Melanerpes uropygialis*). *The Birds of North America Online* (A. Poole, Ed.). Ithaca, New York: Cornell Lab of Ornithology; Accessed April 28, 2011. <http://bna.birds.cornell.edu/bna/species/532>.
- Epps, C. W., P. J. Palsboll, J. D. Wehausen, R. R. Ramey II, and D. R. McCullough. 2005. Highways block gene flow and cause rapid decline in genetic diversity of desert bighorn sheep. *Ecology Letters* 8: 1029-1038.
- Evens, JM and S.L. Hartman. 2007. Vegetation Survey and Classification for the Northern and Eastern Colorado Desert Coordinated Management Plan (NECO). California Native Plant Society. Sacramento, CA.
- Galati and Blek, LLP. 2010. Applicant's ground-based and field-verified delineation of desert dry wash woodland
- Garrett, K., and J. Dunn. 1981. *Birds of Southern California: Status and Distribution*. Los Angeles, California: Los Angeles Audubon Society.
- Gervais, J. A., D. K. Rosenberg, and L. A. Comrack. 2008. Burrowing owl (*Athene cunicularia*). Pages 218-226 in California bird species of special concern: a ranked assessment of species, subspecies, and distinct populations of birds of immediate conservation concern in California (W. D. Shuford and T. Gardali, editors). Western Field ornithologists and California Department of Fish and Game, Studies of Western Birds 1: 1-450.
- Hagerty, B. E., K. E. Nussear, T. C. Esque and C. R. Tracy. 2011. Making molehills out of mountains: landscape genetics of the Mojave desert tortoise. *Landscape Ecology* 26(2): 267-280.
- Harless, M. L., A. D. Walde, D. K. Delaney, L. L. Pater and W. K. Hayes. 2009. Home range, spatial overlap, and burrow use of the desert tortoise in the West Mojave desert. *Copeia*(2): 378-389.
- Harrity, E.J. and C.J. Conway. 2018. Dispersal and Migration Behavior of Yuma Ridgway's rails. Wildlife Research Report #2018-01. Idaho Cooperative Fish and Wildlife Research Unit. Moscow, Idaho.
- Harrity, E.J. and C.J. Conway. 2019. Dispersal and Migration Behavior of Yuma Ridgway's Rails. 2019 Annual Report. Wildlife Research Report #2019-01. Idaho Cooperative Fish and Wildlife Research Unit, Moscow, Idaho.
- Haug, E.A., B.A. Millsap, and M.S. Martell. 1993. Burrowing Owl (*Speotyto cunicularia*). In A. Poole and F. Gill, eds. *The Birds of North America*, No. 61. The Academy of Natural Sciences, Philadelphia, Pennsylvania, and The American Ornithologists' Union, Washington, D.C.
- Hollingsworth, B. D., and K. R. Beaman. 1999. Mojave Fringe-Toed Lizards (*Uma scoparia*). Species Accounts – West Mojave Plan [www.blm.gov/ca/pdfs/cdd\\_pdfs/fringe1.PDF](http://www.blm.gov/ca/pdfs/cdd_pdfs/fringe1.PDF)
- Holland, R.F. 1986. Preliminary Descriptions of the Terrestrial Natural Communities of California. The Resources Agency, Department of Fish and Wildlife, State of California.

- Humple, D. 2008. Loggerhead Shrike (*Lanius ludovicianus*). In: Shuford, W. D., and Gardali, T., eds. California Bird Species of Special Concern: A Ranked Assessment of Species, Subspecies, and Distinct Populations of Birds of Immediate Conservation Concern in California. Studies of Western Birds 1. Western Field Ornithologists, Camarillo, California, and California Department of Fish and Game, Sacramento.
- Ironwood Consulting (Ironwood). 2017. Biological Resource Technical Report – Palen Solar PV Project. Submitted to EDF Renewable Energy. April 12, 2017.
- Ironwood Consulting (Ironwood). 2019. Biological Resource Technical Report – Athos Renewable Energy Project. Submitted to Aspen Environmental Group. January 2019.
- Jarvis, J. M. 2009. The Natural History of the Mojave Fringe-Toed Lizard *Uma scoparia*: The Northern Lineage, Amargosa River, CA. Master's Thesis. California State University, Fullerton.
- Jennings, M. R. and M. P. Hayes. 1994. Amphibian and reptile species of special concern in California. California Department of Fish and Game. Rancho Cordova 255 p.
- Jepson, Willis Linn, and J. C. Hickman. 1993. The Jepson manual: higher plants of California. Berkeley, CA: University of California Press.
- Jones, L.C., and R.E. Lovich. 2009. Lizards of the American Southwest. Tucson, Arizona: Rio Nuevo Publishers. September 30, 2009.
- Kagan, R.A., T.C. Viner, P.W. Trail, and E.O. Espinoza. 2014. Avian mortality at solar energy facilities in southern California: a preliminary analysis. National Fish and Wildlife Forensics Laboratory. Ashland, Oregon.
- Karl, A. 2013a. Summary of Spring Wildlife and Plant Surveys. Letter report to California Energy Commission dated May 16, 2013.
- Kenney, M. 2010. Geomorphic Evaluation of Potentially Affected Aeolian Sand Migration Regions for Reconfigured Alternatives 2 and 3 Associated with the Wind Fence, Palen Solar Power Project (PSPP), Chuckwalla Valley, Riverside County, CA. July 20, 2010.
- Kochert, M.N., K. Steenhof, C.L. McIntyre, and E.H. Craig. 2002. Golden Eagle (*Aquila chrysaetos*). In The Birds of North America, No. 684, edited by A. Poole and F. Gill. The Birds of North America, Inc. Philadelphia.
- Lancaster, J. T., Bedrossian, T. L., and Holland, P., 2014, Eolian System Mapping for the Desert Renewable Energy Conservation Plan, California Geological Survey, 54p., 4 plates (multiple map scales).
- Levenstein, K., A. Chatfield, W. Erickson, and K. Bay. 2014. Fall 2013 Avian Field Surveys for the Palen Solar Electric Generating System, Riverside County, California. Prepared for Palen Solar Holdings, LLC. February 13, 2014.
- Levenstein, K. and C. Nations. 2013. Fall 2013 Nocturnal Migration Surveys for the Palen Solar Electric Generating System, Riverside County, California. Final Report. Prepared for Palen Solar Holdings, LLC. Prepared by Western EcoSystems Technology, Inc. (WEST), Cheyenne, Wyoming.
- MacKay, P. 2003. Mojave Desert wildflowers: a field guide to wildflowers, trees, and shrubs of the Mojave Desert, including the Mojave National Preserve, Death Valley National Park, and Joshua Tree National Park. A Falcon Guide. Guilford, CT: Falcon. 338 p. 65313.
- Mayhew, W. W. 1965. Adaptations of the amphibian, *Scaphiopus couchii* to desert conditions. American Midland Naturalist: pp 95-109.
- Mayhew, W. W. 1965. Reproduction in the sand-dwelling lizard *Uma inornata*. Herpetologica 21, pp. 39-55.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.

- McCreedy, C. 2008. Gila Woodpecker (*Melanerpes uropygialis*). The Desert Bird Conservation Plan. California Partners in Flight. Accessed April 28, 2011. <http://www.prbo.org/calpif/html/docs/desert.html>.
- McGrew, J.C. 1979. *Vulpes macrotis*. Mammalian Species 123:1–6.
- Mills, S.G., J.B. Dunning, Jr., and J.M. Bates. 1989. Effects of Urbanization on Breeding Bird Community Structure in Southwestern Desert Habitats. The Condor 91:416–428.
- Muhs, D.R., Reynolds, R.R., Been, J., Skipp, G., 2003. Eolian sand transport pathways in the southwestern United States: importance of the Colorado River and local sources. Quaternary International, 104, 3-18.
- Munz, Philip A, and David D. Keck, 1973. A California flora and supplement. Berkeley, CA: University of California Press.
- Murphy, R. W., T. L. Trepanier, and D. J. Morafka. 2006. Conservation genetics, evolution and distinct population segments of the Mojave fringe-toed lizard, *Uma scoparia*. Journal of Arid Environments 67:226–247.
- Nussear, K.E., Esque, T.C., Inman, R.D., Gass, Leila, Thomas, K.A., Wallace, C.S.A., Blainey, J.B., Miller, D.M., and Webb, R.H. 2009. Modeling habitat of the desert tortoise (*Gopherus agassizii*) in the Mojave and parts of the Sonoran Deserts of California, Nevada, Utah, and Arizona: U.S. Geological Survey Open-File Report 2009-1102, 18 p.
- O'Connor, M. P., L. C. Zimmerman, D. E. Ruby, S. J. Bulova and J. R. Spotila. 1994. Home range size and movements by desert tortoises, *Gopherus agassizii*, in the eastern Mojave Desert. Herpetological Monographs 8: 60-71.
- Oftedal OT. 2002. Nutritional ecology of the desert tortoise in the Mohave and Sonoran deserts, Chapter 9. In Van Devender TR, editor. The Sonoran Desert Tortoise: Natural History, Biology, and Conservation. Tucson: The University of Arizona Press and Arizona-Sonora Desert Museum, pp. 194–241.
- Oftedal OT, Hillard S, Morafka DJ. 2002. Selective spring foraging by juvenile desert tortoises (*Gopherus agassizii*) in the Mojave Desert: evidence of an adaptive nutritional strategy. Chelonian Conservation Biology 4: 341–352.
- Pagel, J.E., D.M. Whittington, and G.T. Allen. 2010. Interim Golden Eagle inventory and monitoring protocols; and other recommendations. Division of Migratory Bird Management, U.S. Fish and Wildlife Service
- Palen Solar Holdings, LLC. 2013. Final Sand Transport Study for the Palen Solar Electric Generating System. Prepared by Fred L. Nials, Geoarchaeological Consultant to Centerline. July 23, 2013.
- Peterson, C. C. 1996. Ecological energetics of the desert tortoise (*Gopherus agassizii*): Effects of rainfall and drought. Ecology 77(6): 1831-1844.
- Philip Williams and Associates (PWA). 2010. Geomorphic Assessment and Sand Transport Impacts Analysis of the Palen Solar Power Project, Appendix C Biology Report. August 2010.
- Pierson, E.D., and W.E. Rainey. 1998. Western mastiff bat, *Eumops perotis*. In Terrestrial Mammal Species of Special Concern in California, edited by B.C. Bolster. [www.dfg.ca.gov/wildlife/nongame/ssc/docs/mammal/species/17.pdf](http://www.dfg.ca.gov/wildlife/nongame/ssc/docs/mammal/species/17.pdf).
- Potter, C. and J. Weigand, 2016. Analysis of Desert Sand Dune Migration Patterns from Landsat Image Time Series for the Southern California Desert. Journal of Remote Sensing & GIS. May 16, 2016.
- Prescott, B.G. 2005. Le Conte's Thrasher Species Account, West Mojave Plan, Bureau of Land Management. Final environmental impact report and statement for the West Mojave plan: a habitat conservation plan and California desert conservation area plan amendment. Moreno Valley (CA): U.S. Dept. of the Interior, Bureau of Land Management, California Desert District.

- Rosenberg, K.V., S.B. Terill, and G.H. Rosenberg. 1987. Value of Suburban Habitats to Desert Riparian Birds. *Wilson Bulletin* 99(4):642–654.
- Rosenberg, K. V., Ohmart, R. D., Hunter, W. C., and Anderson, B. W. 1991. *Birds of the Lower Colorado River Valley*. Univ. Ariz. Press, Tucson
- Rostal, D. C., V. A. Lance, J. S. Grumbles and A. C. Alberts. 1994. Seasonal reproductive cycle of the desert tortoise (*Gopherus agassizii*) in the eastern Mojave Desert. *Herpetological Monographs* 0(8): 72-82.
- Sawyer, J.O., Jr. and T. Keeler-Wolf. 1995. *A Manual of California Vegetation*. California Native Plant Society, Sacramento, California.
- Sawyer, J.O., Jr., T. Keeler-Wolf, and J. M. Evans. 2009. *A Manual of California Vegetation*. Second edition. California Native Plant Society Press, Sacramento, CA.
- Shuford, W. D., and Gardali, T., editors. 2008. *California Bird Species of Special Concern: A ranked assessment of species, subspecies, and distinct populations of birds of immediate conservation concern in California*. Studies of Western Birds 1. Western Field Ornithologists, Camarillo, California, and California Department of Fish and Game, Sacramento.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey. Available online at <https://websoilsurvey.nrcs.usda.gov/>. Accessed July 2020
- Solar Millennium. 2010a. Spring Survey Protocols, Biological Resources. Palen Solar Power Project. CEC Docket No. 09-AFC-7. April 22 2010.
- Solar Millennium. 2010b. Wildlife Movement and Desert Tortoise Habitat Connectivity. Palen Solar Power Project. CEC Docket No. 09-AFC-7. May 17, 2010.
- Solar Millennium. 2010c. (TN 58454). Palen Solar Power Project golden Eagle Survey Results. Dated September 13, 2010.
- Solar Millennium. 2010d. (TN 58106). Palen Solar Power Project Application for Certification Responses to CEC Information Requests Reconfigured Alternatives 2 and 3 Biological Resources 09-AFC-7. Dated July 21, 2010.
- Stebbins, R. C. 2003. *Western reptiles and amphibians*. Houghton Mifflin Company, New York, New York. 3<sup>rd</sup> ed.
- Tetrattech and A. E. Karl. 2011. *Biological Resources Technical Report*, McCoy Solar Energy Project, Riverside County, CA.
- Turner, F. B., D. C. Weaver and J. C. Rorabaugh. 1984. Effects of reduction in windblown sand on the abundance of the fringe-toed lizard (*Uma inornata*) in the Coachella Valley, California. *Copeia* (2): 370-378.
- U. S. Fish and Wildlife Service (USFWS). 1992. Field Survey Protocol for Any Federal Action that May Occur Within the Range of the Desert Tortoise.
- U. S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened wildlife and plants: Determination of threatened status for the Mojave population of the desert tortoise. USFWS. Ventura, CA. 50 CFR Part 17.
- U. S. Fish and Wildlife Service (USFWS). 1994. Endangered and threatened wildlife and plants: proposed determination of critical habitat for the Mojave population of the desert tortoise. 17: 45748-45768.
- U.S. Fish and Wildlife Service(USFWS). 2009. Desert Tortoise (Mojave Population) Field Manual: (*Gopherus agassizii*). Region 8, Sacramento, California.
- U.S. Fish and Wildlife Service (USFWS). 2010a. Revised pre-project survey protocols for the desert tortoise (*Gopherus agassizii*).

- U. S. Fish and Wildlife Service (USFWS). 2010b. Mojave Population of the Desert Tortoise (*Gopherus agassizii*) 5-Year Review: Summary and Evaluation. U.S. Fish and Wildlife Service. Desert Tortoise Recovery Office. Reno, Nevada. September 30, 2010.
- U. S. Fish and Wildlife Service (USFWS). 2010c. Preparing for any action that may occur within the range of the Mojave desert tortoise (*Gopherus agassizii*). U. S. Fish and Wildlife Service (USFWS). 2011. Revised recovery plan for the Mojave population of the desert tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California. 222 pp.
- U. S. Fish and Wildlife Service (USFWS). 2012. Biological Opinion for the K Road Moapa Solar Project, Moapa River Indian Reservation, Clark County, Nevada. File No. 84320-2011-F-0430 & 1-5-05-FW-536, Tier 5. March 7, 2012.
- U.S. Fish and Wildlife Service (USFWS). 2016. Bald and Golden Eagles: Population demographics and estimation of sustainable take in the United States, 2016 update. Division of Migratory Bird Management, Washington D.C., USA.
- U.S. Fish and Wildlife Service (USFWS). 2019a. Preparing for any action that may occur within the range of the Mojave desert tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service (USFWS) October 8, 2019. 22 pp.
- U.S. Fish and Wildlife Service (USFWS). 2019b. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2018 Annual Report. Report by the Desert Tortoise Recovery office, U.S. Fish and Wildlife Service, Reno, Nevada.
- U.S. Fish and Wildlife Service (USFWS). 2020. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2019 Annual Reporting DRAFT. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- Wehausen, J. D. 2005. Nutrient predictability, birthing seasons, and lamb recruitment for desert bighorn sheep. Pages 37-50 in J. Goerriksen and J. M. Andre, editors. Proceedings of the Sweeney Granite Mountains Desert Research Center 1978-2003: a quarter century of research and teaching. University of California, Riverside, USA.
- Weigand, J. and S. Fitton. 2008. Le Conte's Thrasher (*Toxostoma lecontei*). In The Draft Desert Bird Conservation Plan: a strategy for reversing the decline of desert-associated birds in California. California Partners in Flight: <http://www.prbo.org/calpif/htmldocs/desert.html>
- WEST, Inc. 2016. Draft Bird and Bat Conservation Strategy for the Palen Solar Photovoltaic Project. October 24, 2016.
- Western Bat Working Group (WBWG). 2018. Species Account and status designations. <http://wbwg.org/western-bat-species/>. Accessed August 2018.
- Western Regional Climate Center. 2016. Blythe airport and Eagle Mountain. <http://www.wrcc.dri.edu/>. Accessed July 15, 2016.
- Whiteaker, L.; Henderson, J.; Holmes, R.; Hoover, L.; Leshner, R.; Lippert, J.; Olson, E.; Potash, L.; Seevers J.; Stein M.; Wogen N. 1998. Survey Protocol for Survey and Manage Strategy 2 Vascular Plants. V 2.0.
- Woodbridge, B. 1998. Swainson's Hawk (*Buteo swainsoni*). In The Riparian Bird Conservation Plan: a strategy for reversing the decline of riparian-associated birds in California. California Partners in Flight. [http://www.prbo.org/calpif/htmldocs/riparian\\_v-2.html](http://www.prbo.org/calpif/htmldocs/riparian_v-2.html)
- Yosef, R. 1996. Loggerhead Shrike (*Lanius ludovicianus*). In: The Birds of North America, No. 231 (A. Poole and F. Gill [eds.]). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C.

Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California, in California Wildlife Habitat Relationships System, California Department of Fish and Wildlife, California Interagency Wildlife Task Group.

Zimbelman, JR; Williams, SH; Tchakerian VP. 1995. Sand transport paths in the Mojave Desert, southwestern United States. In: Tchakerian, VP, ed. Desert Aeolian processes. London: Chapman and Hall: pp 101-129.



**Photo 1.** Vegetation community – Sonoran creosote bush scrub



**Photo 2.** Vegetation community – Saltbush scrub





**Photo 3.** Vegetation community – Desert dry wash woodland



**Photo 4.** Active aeolian sand





**Photo 5.** Mohave fringe-toed lizard



**Photo 6.** Canid burrow with burrowing owl white wash



**Photo 7.** Camera station – burro deer



**Photo 8.** Camera station – bob cat





**Photo 9.** Camera station – coyote



**Photo 10.** Camera station - jackrabbit



**Photo 11.** Crucifixion thorn



**Photo 12.** Eriastrum harwoodii






**Photo 13.** *Astragalus harwoodii*



**Photo 14.** *Cryptantha costata*








**FIGURE 1**  
**General Vicinity**  
**Arica Solar**





Ironwood  
Consulting



0 4  
Miles



Arica Solar Project

Ephemeral Drainage

Aqueduct



Intermittent Water Feature




Watershed Boundary

**FIGURE 2**

**Hydrologic Unit Map**

**Arica Solar**







Source: SSURGO 2017

**Ironwood**  
Consulting

 Arica Solar  
Project Boundary

**Soils**


 *Rositas-Dune land-  
Carsitas (s1136)*

 *Vaiva-Quilotosa-Hyder-  
Cipriano-Cherioni (s1141)*


**FIGURE 3**

**Soils**



**Arica Solar**







**Ironwood**  
Consulting

 Ephemeral Drainage  
 Arica Solar Project Boundary




0 1,000 2,000  
Feet

**FIGURE 4**

**Aeolian Sand Transport**

**Arica Solar**








**Ironwood**  
Consulting



0 1,000 2,000  
Feet

**Vegetation Communities**

-  Desert Dry Wash Woodland
-  Saltbush Scrub
-  Sonoran Creosote Bush Scrub



Ephemeral Drainage


Arica Solar Project  
Boundary

**FIGURE 5**

**Vegetation Communities**

**Arica Solar**









**Ironwood**  
Consulting



0 1,000 2,000  
Feet

-  Avian Count Survey Line
-  10-meter Survey Area (Fall 2019)
-  BUOW Surveys #2 - 4 (Spring 2020)
-  20-meter Survey Area (Spring 2020)




Arica Solar Project Boundary

**FIGURE 6**

**Study Area**

**Arica Solar**





Arica Solar  
Boundary

**Ironwood**  
Consulting



0 1,000 2,000  
Feet



Couch's spadefoot toad,  
Potential Habitat



Desert Tortoise Burrow-  
Class 4



Mojave Fringe-toed Lizard,  
Live Individual



Mojave Fringe-toed Lizard-  
Potential Habitat



Mojave Fringe-toed Lizard  
Species Distribution Model


**FIGURE 7**

**Noteworthy Reptile and  
Amphibian Observations**

**Arica Solar**




P:\GIS\Ironwood\Shared\Projects\Arica\Arica\_DT\_TCA\_Linkages.mxd



**FIGURE 8**  
**Desert Tortoise**  
**Conservation Areas (TCAs)**  
**and Linkages**

Arica Solar





**Ironwood  
Consulting**



0 1,000 2,000  
Feet

- Burrowing Owl  
(burrow with sign)
- ▲ Burrowing Owl  
(live individual flushed  
or in flight)
- Burrowing Owl  
(live individual at burrow)
- Leconte's Thrasher  
(live individual)

- ★ Loggerhead Shrike  
(perching or in flight)
- Prairie Falcon  
(in flight)
- ▼ Short-eared Owl  
(live individual)
- ⬠ Swainson's Hawk  
(live individual)


Arica Solar  
Project Boundary

**FIGURE 9**

**Noteworthy Avian  
Observations**

**Arica Solar**





Adapted from: Corvus, 2018.

**Ironwood**  
Consulting



0 2  
Miles

**Season Last Surveyed**

- ▲ 2013-2014
- ▲ 2014-2015



Cliff Nests Monitored During  
BLM Raven Surveys



GOEA Sighting (Date of Sighting)



Arica Solar Project Location




Joshua Tree National Park




Arica Solar Project  
4-mile Buffer

**Number of  
Surveys Conducted**




**FIGURE 10**  
**Golden Eagle**  
**Survey Results**  
**2010-2015**  
**Arica Solar**





**Ironwood**  
Consulting



- Camera Station
- Arica Solar Project Boundary

- Burrow Deer (scat)
- ▲ Burrow Deer (tracks)
- Canid Burrow (inactive)
- ✕ Canid Carcass

- Coyote (burrow - inactive)
- ▼ Coyote (live individual)
- ✕ Coyote (carcass)


- Kit Fox (burrow - inactive)
- ▲ Desert Kit Fox (burrow - active)
- ✕ Desert Kit Fox (carcass)

**FIGURE 11**

**Noteworthy Mammal Observations**

**Arica Solar**





Ironwood  
Consulting




0 2  
Miles

**FIGURE 12**

**Wildlife Connectivity**


**Arica Solar**





**Ironwood**  
Consulting

Arica Solar Project Boundary



**FIGURE 13**

**Noteworthy Plant  
Observations**

**Arica Solar**

## APPENDIX A

### Summary of Survey Results 2019-2020 Arica Solar Project

#### A-1. Daily Survey Personnel

Date	Survey Type	Surveyors	Weather	Average Survey Hours
10/9/2019	Fall Wildlife & Plant	M Adams, M. Baker, M. Basset, K. Black, M. Bratton, S. Clegg, M. Cloud-Hughes, A. Drummer, K. Hayes, T. Hobbs, M. Honer, C. Keaton, D. Kesonie, M. Lavender, A. Layden, W. McBride, C. McClurg, M. Moon, S. Nelson, S. Nielsen, B. Payne, J. St. Pierre, M. Sally, C. Slaughter, A. Walters, Z. Webb, J. Yerger, M. Zhou	58-97°F	10
10/10/2019	Fall Wildlife & Plant	M. Baker, M. Basset, K. Black, M. Bratton, L. Chow, S. Clegg, M. Cloud-Hughes, A. Drummer, M. Dipane, K. Hayes, T. Hobbs, M. Honer, C. Keaton, D. Kesonie, M. Lavender, A. Layden, W. McBride, C. McClurg, M. Moon, S. Nelson, S. Nielsen, B. Payne, J. St. Pierre, M. Sally, C. Slaughter, E. Thorn, A. Walters, Z. Webb, J. Yerger, M. Zhou	62-92°F	10
10/11/2019	Fall Wildlife & Plant	M Adams, M. Baker, M. Basset, K. Black, M. Bratton, S. Clegg, M. Cloud-Hughes, A. Drummer, K. Hayes, T. Hobbs, M. Honer, C. Keaton, D. Kesonie, M. Lavender, A. Layden, W. McBride, C. McClurg, M. Moon, S. Nelson, S. Nielsen, B. Payne, J. St. Pierre, M. Sally, C. Slaughter, A. Walters, Z. Webb, J. Yerger, M. Zhou	61-77°F	8
10/12/2019	Fall Wildlife & Plant	M. Baker, M. Basset, K. Black, M. Bratton, S. Clegg, M. Cloud-Hughes, A. Drummer, K. Hayes, T. Hobbs, M. Honer, C. Keaton, D. Kesonie, M. Lavender, A. Layden, W. McBride, C. McClurg, M. Moon, S. Nelson, S. Nielsen, B. Payne, J. St. Pierre, M. Sally, C. Slaughter, E. Thorn, A. Walters, Z. Webb, J. Yerger, M. Zhou	55-79°F	10
3/21/2020	Spring Plant & Avian	M. Baker, K. Black, L. Chow, M. Cloud-Hughes, D. Kesonie, M. Lavender, W. McBride, S. Nielsen, J. St. Pierre	45-63 °F, wind 5mph	10
3/23/2020	Spring Plant & Avian	M. Baker, L. Chow, M. Cloud-Hughes, D. Kesonie, M. Lavender, W. McBride, S. Nielsen, J. St. Pierre, J. Yerger	54-72 °F , wind 8mph	10

<b>Date</b>	<b>Survey Type</b>	<b>Surveyors</b>	<b>Weather</b>	<b>Average Survey Hours</b>
3/24/2020	Spring Plant & Avian	M. Baker, L. Chow, M. Cloud-Hughes, D. Kesonie, M. Lavender, W. McBride, S. Nielsen, J. St. Pierre, J. Yerger	62-65 °F, wind 5-15mph	10
3/25/2020	Spring Plant & Avian	M. Baker, L. Chow, M. Cloud-Hughes, D. Kesonie, M. Lavender, W. McBride, S. Nielsen, J. St. Pierre, J. Yerger	53-72°F, wind 7mph	10
5/1/2020-5/29/2020	Spring 2020 wildlife cameras at underpasses	L. Chow, Z. Webb, M. Lavender		10
5/7/2020	BUOW Survey #2	L. Chow, Z. Webb	67-102 °F	11
5/8/2020	BUOW Survey #2	L. Chow, Z. Webb	66-101 °F	11
5/21/2020	Jurisdictional Delineation Survey	L. Chow, E. Thorn, D. Kesonie, J. St. Pierre, S. Nielsen	51-83°F	10
6/17/2020	BUOW Survey #3	A. Shaub, M. Lavender	61-96°F	11
6/18/2020	BUOW Survey #3	A. Shaub, M. Lavender	74-96°F	11
7/14/2020	BUOW Survey #4	M. Lavender, Z. Webb	78-112°F	11
7/15/2020	BUOW Survey #4	M. Lavender, Z. Webb	78-112°F	11



## A-2. Summary of Noteworthy Reptile and Amphibian Observations

Reptile/Amphibian Species	Reptile/Amphibian Sign Types	Notes	Date
desert tortoise	Burrow	Class 4, possible tortoise hiding spot, looks like modified rat burrow, by canid.	2019-10-10
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	+5 more juveniles within 200 m, +1 adult within 200 m	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-11
Mojave fringe toed lizard	Live Individual	-	2019-10-12
Mojave fringe toed lizard	Live Individual	-	2019-10-12
Mojave fringe toed lizard	Live Individual	-	2020-03-21
Mojave fringe toed lizard	Live Individual	-	2020-03-21
Mojave fringe toed lizard	Live Individual	-	2020-03-21
Mojave fringe toed lizard	Live Individual	-	2020-03-21
Mojave fringe toed lizard	Live Individual	4 individuals	2020-03-23
Mojave fringe toed lizard	Live Individual	4 individuals	2020-03-24
Mojave fringe toed lizard	Live Individual	-	2020-03-24
Mojave fringe toed lizard	Live Individual	-	2020-03-24
Mojave fringe toed lizard	Live Individual	-	2020-03-24
Mojave fringe toed lizard	Live Individual	-	2020-03-25
Couch's spadefoot toad	Potential Habitat	-	2019-10-09
Couch's spadefoot toad	Potential Habitat	-	2019-10-09

<b>Reptile/Amphibian Species</b>	<b>Reptile/Amphibian Sign Types</b>	<b>Notes</b>	<b>Date</b>
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10
Couch's spadefoot toad	Potential Habitat	-	2019-10-10

### A-3. Summary of Noteworthy Avian Observations

Bird Species	Sign Type	Notes	Date
Burrowing owl	Live individual, burrow. Pellets, whitewash, feathers	Flying, flushed from at hole	2019-10-09
Burrowing owl	Live individual	Flying, Resting	2019-10-10
Burrowing owl	Live individual, burrow. Pellets, whitewash, feathers	Flying	2019-10-10
Burrowing owl	Live individual	Flying	2019-10-11
Burrowing owl	Live individual	Flying, Resting	2019-10-11
Burrowing owl	Live individual	Flying, Resting	2019-10-11
Burrowing owl	Live individual, burrow, feathers	Resting	2019-10-11
Burrowing owl	Live individual, burrow, pellets, tracks	Flying	2019-10-11
Burrowing owl	Live individual, burrow. Pellets, whitewash, feathers, tracks	Flying	2019-10-11
Burrowing owl	Live individual, whitewash, feathers	Resting	2019-10-12
Burrowing owl	Live individual	Flying	2020-03-21
Burrowing owl	Live individual, burrow, Pellets, whitewash, feathers	Flying	2020-03-23
Burrowing owl	Live individual, burrow, whitewash	Resting, Territorial/Combat	2020-03-24
Burrowing owl	Burrow, pellets, whitewash, feathers	-	2019-10-09
Burrowing owl	Burrow, pellets, whitewash, feathers	-	2019-10-10
Burrowing owl	Burrow, feathers	-	2019-10-11
Burrowing owl	Burrow, pellets, whitewash, feathers	Several burrows with sign	2020-03-24
Burrowing owl	Feathers	former canid burrow	2019-10-11
Short-eared owl	Live individual, pellets, whitewash	Flying, roosting	2020-03-23
Leconte's thrasher	Live individual	Singing	2020-03-24
Swainson's hawk	Live individual	Flying	2020-03-25
Prairie falcon	Live individual	Flying	2019-10-09
Prairie falcon	Live individual	Flying low to ground, Foraging	2019-10-10
Prairie falcon	Live individual	Flying	2019-10-10
Prairie falcon	Live individual	Flying	2019-10-10
Prairie falcon	Live individual	Flying	2019-10-11
Prairie falcon	Live individual	Flying, Mating/Courting	2019-10-11
Prairie falcon	Live individual	Flying	2020-03-23
Loggerhead shrike	Live individual	Flying, Perching	2019-10-09
Loggerhead shrike	Live individual	Flying, Perching	2019-10-10
Loggerhead shrike	Live individual	Perching	2019-10-11
Loggerhead shrike	Live individual	Singing	2019-10-14



Bird Species	Sign Type	Notes	Date
Loggerhead shrike	Live individual	Flying, Perching	2020-03-21
Loggerhead shrike	Live individual	Singing	2020-03-23
Loggerhead shrike	Live individual	Perching, Singing, Other	2020-03-23
Loggerhead shrike	Live individual	Foraging	2020-03-23
Loggerhead shrike	Live individual	Flying, Territorial/Combat possible nest nearby	2020-03-23
Loggerhead shrike	Live individual	Singing	2020-03-24

#### A-4. Avian Counts Summary

Avian Species	Date				Species Total
	03/21/20	03/23/20	03/24/20	03/25/20	
American kestrel ( <i>Falco sparverius</i> )			1		1
American pipit ( <i>Anthus rubescens</i> )				1	1
Anna's hummingbird ( <i>Calypte anna</i> )		2			2
Ash-throated flycatcher ( <i>Myiarchus cinerascens</i> )				1	1
Black-tailed gnatcatcher ( <i>Polioptila melanura</i> )		3			3
Black-throated sparrow ( <i>Amphispiza bilineata</i> )			2	2	4
Blue-gray gnatcatcher ( <i>Polioptila caerulea</i> )	1		1	2	4
Cactus wren ( <i>Campylorhynchus brunneicapillus</i> )			1		1
Cliff swallow ( <i>Petrochelidon pyrrhonota</i> )		2	1	2	5
Common raven ( <i>Corvus corax</i> )	2	1	13	2	18
Cooper's hawk ( <i>Accipiter cooperii</i> )			1		1
Gambel's quail ( <i>Callipepla gambelii</i> )			1	3	4
Horned lark ( <i>Eremophila alpestris</i> )	1	1	5		7
House finch ( <i>Haemorhous mexicanus</i> )	1				1
LeConte's thrasher ( <i>Toxostoma lecontei</i> )				1	1
Lesser goldfinch ( <i>Spinus psaltria</i> )		1			1
Lesser nighthawk ( <i>Chordeiles acutipennis</i> )				1	1
Loggerhead shrike ( <i>Lanius ludovicianus</i> )		5	3		8
Mourning dove ( <i>Zenaida macroura</i> )		3		2	5
Northern mockingbird ( <i>Mimus polyglottos</i> )		1	1	1	3
Northern rough-winged swallow ( <i>Stelgidopteryx serripennis</i> )			2		2
Osprey ( <i>Pandion haliaetus</i> )			2		2
Prairie falcon ( <i>Falco mexicanus</i> )			1		1
Red-tailed hawk ( <i>Buteo jamaicensis</i> )			2		2
Tree swallow ( <i>Tachycineta bicolor</i> )		3			3
Turkey vulture ( <i>Cathartes aura</i> )	2	23	66	20	111
Verdin ( <i>Auriparus flaviceps</i> )		5	1		6
Violet-green swallow ( <i>Tachycineta thalassina</i> )			5	1	6
Western meadowlark ( <i>Sturnella neglecta</i> )			1	2	3
Yellow-rumped warbler ( <i>Setophaga coronata</i> )	1		2		3
<b>Total Observed:</b>	<b>8</b>	<b>50</b>	<b>112</b>	<b>41</b>	<b>211</b>

#### A-5. Noteworthy Mammal Observations

Mammal Species	Mammal Sign Types	Active/inactive	Comments	Date
Desert Kit Fox	Burrow	Active	Scat out in the open	2019-10-09
Desert Kit Fox	Burrow	Active	One burrowing owl feather in entrance	2019-10-09
Desert Kit Fox	Burrow	Active	Tracks	2019-10-09
Desert Kit Fox	Burrow	Active	Scat	2019-10-09
Desert Kit Fox	Burrow	Active	Scat	2019-10-10
Desert Kit Fox	Burrow	Active	Scat	2019-10-10
Desert Kit Fox	Burrow	Active	Scat	2019-10-10
Desert Kit Fox	Burrow	Active	Dig marks, tracks	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	Old scat	2019-10-09
Desert Kit Fox	Burrow	Inactive	Dig marks (old)	2019-10-10
Desert Kit Fox	Burrow	Inactive	Old scat	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-10
Desert Kit Fox	Burrow	Inactive	-	2019-10-11
Desert Kit Fox	Burrow	Inactive	-	2019-10-11
Desert Kit Fox	Scat	Inactive	-	2019-10-09
Desert Kit Fox	Burrow	Inactive	Old scat, unknown canid carcass nearby - old and disarticulated	2019-10-10
Desert Kit Fox	Carcass	-	Old kit fox skull	2020-03-24
Canid	Burrow	Inactive	-	2019-10-09
Canid	Burrow	Inactive	-	2019-10-09
Canid	Burrow	Inactive	-	2019-10-10
Canid	Burrow	Inactive	Old scat	2019-10-10
Canid	Burrow	Inactive	-	2019-10-11
Canid	Burrow	Inactive	Burrowing owl whitewash at entrance	2019-10-11
Canid	Burrow	Inactive	Old scat	2019-10-10
Canid	Carcass	Inactive	-	2019-10-11

<b>Mammal Species</b>	<b>Mammal Sign Types</b>	<b>Active/inactive</b>	<b>Comments</b>	<b>Date</b>
Canid	Complex	Inactive	-	2019-10-09
Burro deer	Dig Marks / Tracks	-	-	2019-10-09
Burro deer	Dig Marks / Tracks	-	-	2019-10-09
Burro deer	Dig Marks / Tracks	-	-	2019-10-10
Burro deer	Scat	-	-	2019-10-09
Burro deer	Scat	-	-	2019-10-09
Burro deer	Scat	-	-	2019-10-09
Burro deer	Scat	-	-	2019-10-10
Burro deer	Scat	-	-	2019-10-10
Burro deer	Scat	-	-	2019-10-10
Burro deer	Scat	-	-	2019-10-11
Burro deer	Scat	-	-	2019-10-11
Burro deer	Scat	-	-	2019-10-12
Coyote	Live Individual	Active	Scat 5m from entrance. burrow in berm.	2019-10-10
Coyote	Complex	Active	Live burrowing owl at burrow	2019-10-10
Coyote	Burrow	Active	Large coyote dig with Scat	2019-10-10
Coyote	Burrow	Active	Fresh Scat	2019-10-10
Coyote	Burrow complex	Inactive	-	2019-10-09
Coyote	Burrow	Inactive	-	2019-10-10
Coyote	Burrow	Inactive	-	2019-10-10
Coyote	Burrow	Inactive	-	2019-10-10
Coyote	Burrow complex	Inactive	-	2019-10-10
Coyote	Burrow	Inactive	-	2019-10-10
Coyote	Carcass	Inactive	Partial carcass	2020-03-21

## A-6. Noteworthy Plant Species Observations

Plant Species	Phenology	Comments	Date
<i>Astragalus insularis</i> var. <i>harwoodii</i> (Harwood's milkvetch)	Flower / Fruit	1 individual	2020-03-21
<i>Astragalus insularis</i> var. <i>harwoodii</i> (Harwood's milkvetch)	Plant dried up / Not chlorophytic	3 individuals	2019-10-11
<i>Eriastrum harwoodii</i> (Harwood's wollystar)	Flower only		2020-03-21
<i>Eriastrum harwoodii</i> (Harwood's wollystar)	Flower only	40 individuals	2020-03-24
<i>Eriastrum harwoodii</i> (Harwood's wollystar)	Flower only	60 individuals	2020-03-24
<i>Eriastrum harwoodii</i> (Harwood's wollystar)	Flower only	8 individuals	2020-03-24
<i>Eriastrum harwoodii</i> (Harwood's wollystar)	Flower only	70	2020-03-25
<i>Castela emoryi</i> (Emory's crucifixion thorn)	Fruit only	Four or five individuals in close proximity	2019-10-10
<i>Castela emoryi</i> (Emory's crucifixion thorn)	Vegetative	One declining individual	2019-10-10
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	70 individuals, some on project, some in buffer	2020-03-24
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	15	2020-03-24
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	4	2020-03-24
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	5000+	2020-03-24
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	estimated 3200+	2020-03-25
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	100+	2020-03-25
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	estimated 3200	2020-03-25
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	100+	2020-03-25
<i>Cryptantha costata</i> (ribbed cryptantha)	Flower / Fruit	Estimated 68	2020-03-25
<i>Cryptantha costata</i> (ribbed cryptantha)	Plant dried up / Not chlorophytic	-	2019-10-11
<i>Mammillaria tetrancistra</i> (fishhook cactus)	Vegetative	-	2019-10-09
<i>Mammillaria tetrancistra</i> (fishhook cactus)	Vegetative	-	2019-10-10
<i>Mammillaria tetrancistra</i> (fishhook cactus)	Vegetative	-	2020-03-18
<i>Mammillaria tetrancistra</i> (fishhook cactus)	Vegetative	-	2020-03-23
<i>Fouquieria splendens</i> (ocotillo)	Vegetative	2 individuals 15 m apart	2019-10-12

## A-7. Invasive Plant Species Observations

Invasive Species	Phenology	Comments	Date
<i>Brassica tournefortii</i> (Sahara mustard)	Flower / Fruit	-	2020-03-21
<i>Brassica tournefortii</i> (Sahara mustard)	Flower / Fruit	-	2020-03-23
<i>Brassica tournefortii</i> (Sahara mustard)	Flower only	-	2020-03-23
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Fruit only	Low density	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	10 individuals	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	along berm	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	Low density	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	1 individual	2019-10-09
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	7 individuals	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	1 individual	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	2 individuals	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	8 individuals	2019-10-10
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	1 individual	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	-	2019-10-11

Invasive Species	Phenology	Comments	Date
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	hotspot	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	3 individuals	2019-10-11
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	approximately 80 individuals	2019-10-12
<i>Brassica tournefortii</i> (Sahara mustard)	Plant dried up / Not chlorophytic	approximately 80 individuals	2019-10-12
<i>Salsola tragus</i> (Russian thistle)	Flower / Fruit	about 100 individuals	2020-03-25
<i>Salsola tragus</i> (Russian thistle)	Flower only	-	2020-03-24
<i>Salsola tragus</i> (Russian thistle)	Vegetative	-	2020-03-21
<i>Salsola tragus</i> (Russian thistle)	Vegetative	-	2020-03-21
<i>Salsola tragus</i> (Russian thistle)	Vegetative	-	2020-03-24
<i>Salsola tragus</i> (Russian thistle)	Vegetative	-	2020-03-24
<i>Sisymbrium irio</i> (tumble mustard)	Flower / Fruit	-	2020-03-21
<i>Sisymbrium irio</i> (tumble mustard)	Flower / Fruit	-	2020-03-23
<i>Sonchus arvensis</i> (sow thistle)	Flower only	-	2020-03-24
<i>Sonchus oleraceus</i> (spiny sowthistle)	Flower / Fruit	-	2020-03-23

## APPENDIX B

### Potential for Special Status Wildlife Species to Occur Arica Solar Project

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>AMPHIBIAN and REPTILES</b>			
<b>Couch's spadefoot toad</b>  <i>Scaphiopus couchii</i>	<p>Occurs along desert washes, desert riparian, palm oasis, desert succulent shrub, and desert scrub habitats. Also found in cultivated cropland areas. Breeds in temporary pools within rocky streambeds, washes, agricultural fields, in road depressions, railroad tracks, and cattle tanks. Pools of water must persist 7 to 8 days to facilitate eggs hatching and larvae transformation</p>	<p>Federal: None</p> <p>State: SSC</p> <p>BLM sensitive</p>	<p>Low to moderate</p> <p>Not observed</p> <p>Potential to occur near water accumulation areas; unconfirmed – no sufficient levels of warm season rain to date</p>
<b>Agassiz's desert tortoise</b>  <i>Gopherus agassizii</i>	<p>higher populations in creosote bush communities with friable soils for burrow construction, with extensive annual blooms, but found in almost every desert habitat</p>	<p>Federal: FT</p> <p>State: ST</p> <p>State: ST</p>	<p>Moderate</p> <p>One class 4 burrow observed near western boundary</p>
<b>Mojave fringe-toed lizard</b>  <i>Uma scoparia</i>	<p>Restricted to fine, loose, wind-blown deposits in sand dunes, dry lakebeds, riverbanks, desert washes, sparse alkali scrub and desert shrub habitats</p>	<p>Federal: None</p> <p>State: SSC</p> <p>BLM sensitive</p>	<p>High</p> <p>Observed</p> <p>Several live individuals in aeolian sandy areas</p>



SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>BIRDS</b>			
<b>Golden eagle</b>  <b>(Nesting and wintering)</b>  <i>Aquila chrysaetos</i>	Typically rolling foothills, mountain areas, sage-juniper flats, desert. Nests on cliffs of all heights and in large trees in open areas. Rugged, open habitats with canyons and escarpments used most frequently for nesting.	Federal: BCC  State: CFP, WL  BLM sensitive	Nesting - lacks suitable habitat  Foraging year-round – low-moderate (uncommon) Not observed
<b>Short-eared owl (Nesting)</b>  <i>Asio flammeus</i>	Year-round residents in N. California and other parts of CA during wintering. Require open country that supports small mammals that also provides adequate vegetation to provide cover for nests includes salt- and freshwater marshes, irrigated alfalfa or grain fields, and ungrazed grasslands and old pastures.	Federal: None  State: SSC	Nesting - lacks suitable habitat  migration or seasonal foraging - moderate  1 individual observed
<b>Western burrowing owl</b>  <i>Athene cunicularia hypugaea</i>	A yearlong resident of open, dry grassland and desert habitats. Uses rodent or other burrows for roosting and nesting cover. In the Colorado Desert, generally occur at low densities in scattered populations	Federal: BCC  State: SSC  BLM sensitive	Nesting – moderate to high  Foraging - high  Live individuals and sign observed
<b>Redhead (Nesting)</b>  <i>Aythya americana</i>	During breeding season may be found along the Colorado River and Salton Sea. Breeds locally in the Central Valley, coastal Southern California, eastern Kern County, and the Salton Sea. Nests in fresh water emergent wetland bordering open water.	Federal: None  State: SSC	Nesting - lacks suitable habitat  Potential migration flyover-moderate  Distant from nearest records - nearest breeding habitat in Salton Sea, not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Ferruginous hawk (Wintering)</b>  <i>Buteo regalis</i>	Most common in grassland and agricultural areas in the southwest. Found in open terrain from grasslands to deserts and are usually associated with concentrations of small mammals.	Federal: BCC  State: WL	Nesting – lacks suitable habitat and outside range  Wintering/migration - moderate  Not observed
<b>Swainson's hawk</b>  <i>Buteo swainsoni</i>	Require large areas of open landscape for foraging, including grasslands and agricultural lands that provide low-growing vegetation for hunting and high rodent prey populations. Typically nest in large native trees such as valley oak, cottonwood, walnut, willow, and occasionally in nonnative trees within riparian woodlands, roadside trees, trees along field borders, isolated trees, small groves, and on the edges of remnant oak woodlands	Federal: BCC  State: ST	Nesting – lacks suitable habitat, outside range  Foraging – moderate, may forage or flyover Project site during migration  Observed in flight
<b>Costa's hummingbird (Nesting)</b>  <i>Calypte costae</i>	Primary habitats are desert wash, edges of desert riparian and valley foothill riparian	Federal: BCC  State: None	Nesting – low, marginal habitat  Foraging – moderate  Not observed
<b>Vaux's swift (Nesting)</b>  <i>Chaetura vauxi</i>	Not known to breed in Riverside or Southern California. They prefer to nest in the hollows inside of large old conifer trees, especially snags, which are entirely lacking from the Project site.	Federal: None  State: SSC	Nesting – lacks suitable habitat  Migration – moderate, potential flyover  not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Mountain plover (Wintering)</b>  <i>Charadrius montanus</i>	habitat includes short-grass prairie or their equivalents, and in southern California deserts are associated primarily with agricultural areas	Federal: BCC  State: SSC  BLM sensitive	Nesting – lacks habitat, outside range  Wintering – not expected  Not observed
<b>Black tern</b>  <i>Chlidonias niger</i>	restricted to freshwater habitats while breeding, can be fairly common on bays, salt ponds, river mouths, and pelagic waters in spring and fall migration (Grinnell and Miller 1944, Cogswell, 1977)	Federal: None  State: SSC	Nesting – lacks habitat  Migration – low, uncommon  Not observed
<b>Northern harrier (Nesting)</b>  <i>Circus cyaneus</i>	Does not commonly breed in desert regions of California, where suitable habitat is limited, but winters broadly throughout California in areas with suitable habitat. Northern harriers forage in open habitats including deserts, pasturelands, grasslands, and old fields.	Federal: None  State: SSC	Nesting – lacks habitat  Foraging/migration - moderate  Observed in flight
<b>Western yellow-billed cuckoo</b>  <i>Coccyzus americanus occidentalis</i>	Breeds along the major river valleys in southern and western New Mexico, and central and southern Arizona. In California, the western yellow-billed cuckoo's breeding distribution is now thought to be restricted to isolated sites in the Sacramento, Amargosa, Kern, Santa Ana, and Colorado River valleys.	Federal: FT, BCC  State: SE  BLM sensitive	Nesting – lacks habitat, outside range  Foraging/migration – low, uncommon  Not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Gilded flicker</b>  <i>Colaptes chrysoides</i>	Stands of giant cactus, Joshua tree, and riparian groves of cottonwoods and tree willows in warm desert lowlands and foothills. Nests primarily in cactus, but also will use cottonwoods and willows of riparian woodlands. May be nearly extinct in California.	Federal: BCC  State: SE  BLM sensitive	Nesting – low, marginal habitat  Migration/foraging – low marginal habitat  Not observed, distant from nearest record
<b>Black swift (Nesting)</b>  <i>Cypseloides niger</i>	Nests in moist crevice or cave on sea cliffs or above the surf, or on cliffs behind, or adjacent to, waterfalls in deep canyons. Forages widely over many habitats.	Federal: BCC  State: SSC	Nesting – lacks suitable habitat  Migration – low, uncommon  Not observed
<b>Willow flycatcher (Nesting)</b>  <i>Empidonax traillii</i>  <b>Southwestern willow flycatcher</b>  <i>E. t. extimus</i>	Most often occurs in broad, open river valleys or large mountain meadows with lush growth of shrubby willows (Serena 1982). Common spring (mid-May to early June) and fall (mid- August to early September) migrant at lower elevations, primarily in riparian habitats throughout the state exclusive of the North Coast.	Federal: None  State: SE  Federal: FE  State: SE	Nesting/wintering- lacks suitable habitat  Migration – low, uncommon Not observed  Nesting/wintering – lacks suitable habitat  Migration – low, uncommon migrant Not observed
<b>California horned lark</b>  <i>Eremophila alpestris actia</i>	A common to abundant resident in a variety of open habitats, usually where trees and large shrubs are absent. Found from grasslands along the coast and deserts near sea level to alpine dwarf-shrub habitat above treeline. In winter, flocks in desert lowlands and other areas augmented by winter visitants, many migrating from outside the state (Garrett and Dunn 1981).	Federal: None  State: WL	Nesting/wintering – high, suitable habitat  Foraging - high  Observed throughout Project site

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Prairie falcon (Nesting)</b>  <i>Falco mexicanus</i>	Occurs in annual grasslands to alpine meadows, but associated primarily with perennial grasslands, savannahs, rangeland, some agricultural fields, and desert scrub. Typically nests at cliffs and bluffs	Federal: BCC  State: WL	Nesting – lacks habitat  Foraging - high  Observed in flight
<b>American peregrine falcon (Nesting)</b>  <i>Falco peregrinus anatum</i>	Rare in the arid southeast, occur and are suspected to breed in the lower Colorado River Valley. Peregrine falcons require open habitat for foraging and prefer breeding sites near water. Nesting habitat includes cliffs, steep banks, dunes, mounds, and some human-made structures	Federal: BCC  State: CFP	Nesting – lacks habitat  Foraging/migration- moderate  Not observed
<b>Sandhill crane (Wintering)</b>  <i>Grus canadensis</i>	Breeds in open wetland habitats surrounded by shrubs or trees. They nest in marshes, bogs, wet meadows, prairies, burned-over aspen stands, and other moist habitats, preferring those with standing water. Outside of known wintering grounds, extremely rare except during migration over much of interior California.	Federal: None  State: SSC	Nesting/wintering/foraging – lacks habitat, outside range  Migration – moderate (flyover)  Not observed
<b>Yellow-breasted chat (Nesting)</b>  <i>Icteria virens</i>	This species occupies shrubby riparian habitat with an open canopy, and will nest in non- native species, including tamarisk.	Federal: None  State: SSC	Nesting - lacks habitat  Migration – moderate (flyover)  Not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Loggerhead shrike (Nesting)</b>  <i>Lanius ludovicianus</i>	Open habitats with scattered shrubs, trees, posts, fences, utility lines, or other perches. Highest density occurs in open-canopied valley foothill hardwood, valley foothill hardwood-conifer, valley foothill riparian, pinyon-juniper, juniper, desert riparian, and Joshua tree habitats	Federal: BCC  State: SSC	Nesting/foraging - high  Observed throughout Project site
<b>Gila woodpecker</b>  <i>Melanerpes uropygialis</i>	In California, this species is found primarily along the Colorado River and in small numbers in Imperial County. In southeastern California, Gila woodpeckers formerly were associated with desert washes extending up to 1 mile from the Colorado River; however, their range may be expanding	Federal: BCC  State: SE  BLM sensitive	Nesting – low, marginal habitat  Foraging – low, marginal habitat  Not observed
<b>Elf owl</b>  <i>Micrathene whitneyi</i>	A very rarely seen spring and summer resident of the Colorado River Valley. Nests in desert riparian habitat with cottonwood, sycamore, willow or mesquite; absent from desert riparian habitat dominated by saltcedar	Federal: BCC  State: SE  BLM sensitive	Nesting – low,  Foraging – low, marginal habitat  Not observed
<b>Long-billed curlew (Nesting)</b>  <i>Numenius americanus</i>	Preferred breeding and winter habitats include large coastal estuaries, upland herbaceous areas, and croplands. On estuaries, feeding occurs mostly on intertidal mudflats.	Federal: BCC  State: WL	Nesting/foraging – lacks suitable habitat  Migration – moderate, flyover  Not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Lucy's warbler (Nesting)</b>  <i>Oreothlypis luciae</i>	An uncommon to common, summer resident and breeder along the Colorado River, common locally in a few other desert areas, and rare near Salton Sea. It occurs in desert typical nesting habitat, mesquite wash and desert riparian habitats, may use abandoned verdin nests	Federal: BCC  State: SSC  BLM sensitive	Nesting – low, marginal habitat  Foraging – low, marginal habitat  Not observed
<b>American white pelican (Nesting colony)</b>  <i>Pelecanus erythrorhynchos</i>	Common spring and fall migrant at Salton Sea and Colorado River. Migrant flocks pass overhead almost any month, but mainly in spring and fall throughout the state, especially in southern California (Cogswell 1977, McCaskie et al. 1979, Garrett and Dunn 1981)	Federal: None  State: SSC	Nesting/wintering/foraging – lacks suitable habitat Migration - moderate, overflight  Not observed
<b>Black-tailed gnatcatcher</b>  <i>Polioptila melanura</i>	A year-round resident in southwestern U.S. and central and northern Mexico, in California, is found in the southeast desert wash habitat from Palm Springs and Joshua Tree National Park south, and along the Colorado River. It is now rare in eastern Mojave Desert north to the Amargosa River, Inyo County. This species nests primarily in wooded desert wash habitat, but also occurs in creosote scrub habitat during the non-breeding season.	Federal: None  State: WL	Nesting/foraging – moderate to high occurrence, suitable habitat present  Observed throughout Project site
<b>Vesper sparrow</b>  <i>Pooecetes gramineus</i>	Fairly common locally in southern deserts in the winter and during migration. Occupies grasslands, croplands, and open brushlands.	Federal: None  State: SSC	Nesting/wintering – lacks suitable habitat  Migration - moderate  Not observed



SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Purple martin</b>  <i>Progne subis</i>	<p>The historical breeding range of the purple martin includes southern California, though populations have shrunk dramatically and neither includes the Colorado Desert. Habitat requirements include adequate nest sites and availability of large aerial insects, and therefore are most abundant near wetlands and other water sources.</p>	<p>Federal: None</p>           <p>State: SSC</p>	<p>Nesting/wintering – lacks suitable habitat</p>  <p>Migration - moderate, flyover</p>           <p>Not observed</p>
<b>Vermilion flycatcher (Nesting)</b>  <i>Pyrocephalus rubinus</i>	<p>They are usually found near water in arid scrub, farmlands, parks, golf courses, desert, savanna, cultivated lands, and riparian woodlands; nesting substrate includes cottonwood, willow, and mesquite.</p>	<p>Federal: None</p>           <p>State: SSC</p>	<p>Nesting/wintering – lacks suitable habitat</p>           <p>Migration/foraging- moderate</p>           <p>not observed</p>
<b>Ridgway's (Yuma) clapper rail</b>  <i>Rallus obsoletus yumanensis</i>	<p>Occurs in inland areas in the southwestern United States. This subspecies is partially migratory, with many birds wintering in brackish marshes along the Gulf of California. Some remain on their breeding grounds throughout the year; for example, the Salton Sea (south) Christmas Bird Count frequently records this species in the fresh-water marshes in and around the Imperial Wildlife Area (Wister Unit). Nesting and foraging habitat occurs only along the Lower Colorado River (from Topock Marsh southward) and around the Salton Sea</p>	<p>Federal: FE</p>           <p>State: ST, CFP</p>	<p>Nesting/wintering – lacks suitable habitat</p>  <p>Migration / dispersal – low, rare (overflight)</p>           <p>Not observed</p>

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Bank swallow (Nesting)</b>  <i>Riparia riparia</i>	A neotropical migrant found primarily in riparian and other lowland habitats in California west of the deserts during the spring-fall period. Uses holes dug in cliffs and river banks for cover. Will also roost on logs, shoreline vegetation, and telephone wires.	Federal: None  State: ST  BLM sensitive	Nesting/wintering – lacks suitable habitat  Migration – moderate, flyover  Not observed
<b>Sonora Yellow warbler (Nesting)</b>  <i>Setophaga petechia sonorana</i>	In southeastern California, this species is known only from the lower Colorado River Valley from the middle of San Bernardino County through Riverside and Imperial Counties. This species commonly uses wet, deciduous thickets for breeding, and seeks a variety of wooded, scrubby habitats in winter	Federal: BCC  State: SSC	Nesting/wintering – lacks suitable habitat  Migration – moderate  Not observed - migration
<b>Lawrence's goldfinch (Nesting)</b>  <i>Spinus lawrencei</i>	Highly erratic and localized in occurrence. Rather common along western edge of southern deserts. Breeds in open oak or other arid woodland and chaparral, near water. Typical habitats in southern California include desert riparian, palm oasis, pinyon-juniper, and lower montane habitats.	Federal: BCC  State: none	Nesting/wintering - low, marginal habitat  Migration moderate  Not observed
<b>Bendire's thrasher</b>  <i>Toxostoma bendirei</i>	Favors open grassland, shrubland, or woodland with scattered shrubs, primarily in areas that contain large cholla, Joshua tree, Spanish bayonet, Mojave yucca, palo verde, mesquite, catclaw, desert-thorn, or agave.	Federal: BCC  State: SSC  BLM sensitive	Nesting – low, marginal habitat  Foraging – moderate  Not observed
<b>Crissal thrasher</b>  <i>Toxostoma crissale</i>	This species prefers habitats characterized by dense, low scrubby vegetation, which, at lower elevations, includes desert and foothill scrub and riparian brush.	Federal: None  State: SSC	Nesting/wintering - low, marginal habitat  Migration moderate  Not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Le Conte's thrasher</b>  <i>Toxostoma lecontei</i>	Occurs primarily in open desert wash, desert scrub, alkali desert scrub, and desert succulent shrub habitats; also occurs in Joshua tree habitat with scattered shrubs.	Federal: None  State: SSC	Nesting/foraging high  Observed
<b>Arizona Bell's vireo</b>  <i>Vireo bellii arizonae</i>  <b>Least Bell's vireo</b>  <i>V. b. pusillus</i>	Subspecies <i>V. b. pusillus</i> (endemic to California and Baja California - state and federally listed) and <i>V.b arizonae</i> are state listed. Bell's vireo is a rare, local, summer resident below about 600 m (2000 ft) in willows and other low, dense valley foothill riparian habitat and lower portions of canyons mostly in San Benito and Monterey Co.; in coastal southern California from Santa Barbara Co. south; and along the western edge of the deserts in desert riparian habitat.	Federal: BCC  State: SE BLM sensitive  Federal: FE  State: SE	Nesting/wintering – lacks suitable habitat Migration moderate  Not observed  Nesting/wintering – lacks suitable habitat Migration moderate  Not observed
<b>Yellow-headed blackbird (Nesting)</b>  <i>Xanthocephalus xanthocephalus</i>	Nests in fresh emergent wetland with dense vegetation and deep water, often along borders of lakes or ponds. Forages in emergent wetland and moist, open areas, especially cropland and muddy shores of lacustrine habitat. Occurs as a migrant and local breeder in deserts	Federal: None  State: SSC	Nesting/wintering/foraging – lacks suitable habitat  Migration – moderate, flyover  Not observed

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>MAMMALS</b>			
<b>Burro deer</b> <i>Odocoileus hemionus eremicus</i>	Occur in early to intermediate successional stages of most forest, woodland, and brush habitats. Prefer a mosaic of various-aged vegetation that provides woody cover, meadow and shrubby openings, and free water	Federal: None State: CPGS	High – sign observed throughout Project site
<b>Desert bighorn sheep</b> <i>Ovis canadensis nelsoni</i>	Habitats used include alpine dwarf-shrub, low sage, sagebrush, bitterbrush, pinyon-juniper, palm oasis, desert riparian, desert succulent shrub, desert scrub, subalpine conifer, perennial grassland, montane chaparral, montane riparian (DeForge 1980, Monson and Sumner 1980, Wehausen 1980). Use rocky, steep terrain for escape and bedding. Remain near rugged terrain while feeding in open habitat	Federal: BLMS State: CFP	Low - unsuitable habitat except as infrequent dispersal between mountain ranges  Not observed
<b>Yuma mountain lion</b> <i>Puma concolor browni</i>	Primarily inhabit the low mountains and extensive wash systems in and around Chuckwalla Bench, Chuckwalla Mountains, Chocolate Mountains, Picacho Mountains, Milpitas Wash, Vinagre Wash, and other washes in that area. Mountain lions typically occur in habitat areas with extensive, well- developed riparian or shrubby vegetation interspersed with irregular terrain, rocky outcrops, and community edges. Restricted to the southern Colorado Desert from Joshua Tree National Park south and east to the Colorado River.	Federal: None State: SSC	Low – marginal habitat  Not observed
<b>American badger</b> <i>Taxidea taxus</i>	Suitable habitat for badgers is characterized by herbaceous, shrub, and open stages of most habitats with dry, friable soils.	Federal: None State: SSC	Moderate to high  Potential sign observed on site

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Desert kit fox</b>  <i>Vulpes macrotis arsipus</i>	Lives in annual grasslands or grassy open vegetation dominated by scattered brush, shrubs, and scrub. Cover provided by occur. Active dens/complexes with sign observed. dens they dig in open, level areas with loose-textured, sandy and loamy soils.	Federal: None  State: CPF	High  Burrows, complexes, scat observed
<b>BATS</b>			
<b>Pallid bat</b>  <i>Antrozous pallidus</i>	Inhabit low elevation (less than 6,000 feet) rocky, arid deserts and canyonlands. Typical roosting habitat is not shrub/steppe grasslands. Day and night roosts include crevices in rocky outcrops and cliffs, however, roosting opportunities may exist outside caves, mines, trees with exfoliating bark, and various human structures (WBWG, 2005)	Federal: None  State: SSC  BLM sensitive	Foraging moderate, roosting low  Not observed  Potentially forage throughout Project
<b>Townsend's big-eared bat</b>  <i>Corynorhinus townsendii</i>	Habitat associations include coniferous forests, deserts, native prairies, riparian communities, active agricultural areas, and coastal habitat types. Foraging associations include edge habitats along streams, adjacent to and within a variety of wooded habitats.	Federal: None  State: SSC  BLM sensitive	Foraging moderate, roosting low-moderate  Not observed  Potentially forage throughout Project
<b>Big brown bat</b>  <i>Eptesicus fuscus</i>	widespread and abundant species has been recorded in virtually every North American vegetation type. Uncommon in hot desert habitats and is absent only from the highest alpine meadows and talus slopes. Vagrant individuals may be seen in any habitat. Uses buildings and other human-made structures for roosting to such an extent that natural roosting habits are under documented	Federal: None  State: none	Low  Not observed  Distant from nearest records

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Spotted bat</b>  <i>Euderma maculatum</i>	Arid, low desert habitats to high elevation conifer forests and prominent rock features appear to be a necessary feature for roosting	Federal: None  State: SSC  BLM sensitive	Low  Not observed  Distant from nearest records
<b>Western mastiff bat</b>  <i>Eumops perotis</i>	Variety of habitats, from desert scrub to chaparral to oak woodland and into the ponderosa pine belt and high elevation meadows of mixed conifer forests	Federal: None  State: SSC  BLM sensitive	Foraging moderate, roosting low  Not observed
<b>Hoary bat</b>  <i>Lasiurus cinereus</i>	Highly associated with forested habitats. Usually are located at the edge of a clearing, although more unusual roosting sites have been reported in caves, beneath rock ledges, woodpecker holes, squirrel nests, building sides, and in dried palm fronds on palm trees.	Federal: None  State: None	Foraging moderate, roosting low  Not observed
<b>Western yellow bat</b>  <i>Lasiurus xanthinus</i>	Recorded below 600 m (2000 ft) in valley foothill riparian, desert riparian, desert wash. This species occurs year-round in California.	Federal: None  State: SSC	Foraging and roosting moderate  Not observed
<b>California leaf-nosed bat</b>  <i>Macrotus californicus</i>	species depends on either caves or mines for roosting habitat. All major maternity, mating, and overwintering sites are in mines or caves (BLM CDD, 2002). California leaf-nosed bat forage almost exclusively among desert wash vegetation within 10 km of their roost (WBWG, 2005)	Federal: None  State: SSC  BLM sensitive	Foraging moderate, roosting low  Not observed  Potentially forage in desert washes



SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Arizona myotis</b>  <i>Myotis occultus</i>	Commonly known from conifer forests from 6,000 to 9,000 feet in elevation, although maternity roosts are known from much lower elevations including areas along the Colorado River in California.	Federal: None  State: SSC	Low Not observed  Distant from nearest records in Blythe
<b>Cave myotis</b>  <i>Myotis velifer</i>	Found primarily at lower elevations of the arid southwest in areas dominated by creosote bush, palo verde, and cactus. This species is a “cave dweller” and caves are the main roosts although this species may also use mines, buildings, and bridges for roosts	Federal: None  State: SSC  BLM sensitive	Low  Not observed Distant from nearest records in Mule Mountains
<b>Yuma myotis</b>  <i>Myotis yumanensis</i>	Associated with permanent sources of water, typically rivers and streams, feeding primarily on aquatic emergent insects. Also use tinajas (small pools in bedrock) in the arid west. Occurs in a variety of habitats including riparian, arid scrublands and deserts, and forests. Roosts in bridges, buildings, cliff crevices, caves, mines, and trees.	Federal: None  State: None  BLM sensitive	Low  Not observed Distant from nearest records in Blythe
<b>Pocketed free-tailed bat</b>  <i>Nyctinomops femorosaccus</i>	Known to occur in the desert from Mar-Aug, when they then migrate out of the area. In California, found primarily in creosote bush and chaparral habitats in proximity to granite boulders, cliffs, or rocky canyons.	Federal: None  State: SSC	Low Not observed  Distant from nearest records in Orocopia Mountains

SPECIES	HABITAT REQUIREMENTS	CONSERVATION STATUS	POTENTIAL TO OCCUR ON PROJECT SITE
<b>Big free-tailed bat</b>  <i>Nyctinomops macrotis</i>	Found generally sea level to 8,000 feet in elevation. This species occurs in desert shrub, . It roosts mostly in the crevices of rocks although may roost in buildings, caves, and tree cavities	Federal: None  State: SSC	Foraging moderate, roosting low  Not observed

Conservation Status

Federal    FE = Federally listed endangered: species in danger of extinction throughout a significant portion of its range  
               FT = Federally listed, threatened: species likely to become endangered within the foreseeable future  
               FCT = Proposed for federal listing as a threatened species  
               BCC = Fish and Wildlife Service: Birds of Conservation Concern:  
               BLMS = Bureau of Land Management Sensitive

State        SSC = State Species of Special Concern  
               CFP = California Fully Protected  
               SE = State listed as endangered  
               ST = State listed as threatened  
               WL = State watch list  
               CPF = California Protected Furbearing Mammal  
               CPGS = California Protected Game Species

\*\* Species not detected during previous surveys may have the potential to occur on the Project site in the future.

## APPENDIX C

### Potential for Special Status Plant Species to Occur Oberon Solar Project

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Chaparral sand verbena <i>Abronia villosa</i> var. <i>aurita</i>	Annual herb; sandy – chaparral, coastal scrub, desert dunes; Imperial, Los Angeles, Orange, Riverside, San Bernardino, San Diego, Ventura; Palen sand dunes, Desert Lily Sanctuary	Federal: none  CRPR: 1B.1  BLM sensitive	75 - 1600	Jan-Sep	Moderate  Not observed
Angel trumpets <i>Acleisanthes longiflora</i>	Perennial herb; sonoran desert scrub (carbonate); known in CA only from one occurrence in the Maria Mountains	Federal: none  CRPR: 2B.3	90 - 95	May	Low – distant from known records  not observed
Desert sand parsley <i>Ammoselinum giganteum</i> / <i>Spermolepis gigantea</i>	Annual herb; Sonoran Desert scrub, Riverside- known in CA only from Hayfields Dry Lake	Federal: none  CRPR: 2B.1	~152	Mar-Apr	Low - distant from known records  Not observed
Small-flowered androstephium <i>Androstephium breviflorum</i>	perennial bulbiferous herb; desert dunes, Mojavean desert scrub (bajada); San Bernardino, Riverside, Inyo; Eastern edge of Eagle Mountains	Federal: none  CRPR: 2B.2	220 - 800	Mar-Apr	Moderate  Not observed

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Harwood's milkvetch <i>Astragalus insularis</i> var. <i>harwoodii</i>	Annual herb; sandy or gravelly - desert dunes, Mojavean Desert scrub; Riverside, San Bernardino, San Diego, Inyo	Federal: none CRPR: 2B.2	0-710	Jan-May	<b>Present</b>
Coachella Valley milkvetch <i>Astragalus lentiginosus</i> var. <i>cochellae</i>	Annual/perennial herb; Desert dunes - Sonoran desert scrub (sandy); endemic to Coachella Valley	Federal: FE CRPR: 1B.2 BLM sensitive	40-655	Feb-May	Low - outside range Not observed
California ayenia <i>Ayenia compacta</i>	Perennial herb; Mojavean desert scrub Sonoran desert scrub; Riverside, San Bernardino, San Diego; Chuckwalla Mountains	Federal: none CRPR: 2B.3	150-1095	Mar-Apr	Low – marginal habitat Not observed
Pink fairy duster <i>Calliandra eriophylla</i>	perennial deciduous shrub Sonoran Desert scrub (sandy or rocky); Imperial, Riverside, San Diego; south of Ford Dry Lake	Federal: none CRPR: 2B.3	120 - 1500	Jan-Mar	Low – marginal habitat Not observed
Sand evening-primrose <i>Chylisimia [Camissonia] arenaria</i>	annual / perennial herb; Sonoran Desert scrub (sandy or rocky); Imperial, Riverside, San Bernardino; Hayfield Lake and Orocopia Mountains	Federal: none CRPR: 2B.2	70-915	Nov-May	Moderate Not observed
Crucifixion thorn <i>Castela emoryi</i>	Perennial deciduous shrub; gravelly - Mojavean desert scrub, Playas, Sonoran Desert scrub, Imperial, Inyo, Riverside, San Bernardino	Federal: none CRPR: 2B.2	90-725	Apr-Oct	<b>Present</b>

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Abram's spurge <i>Chamaesyce abramsiana</i>	Annual herb; sandy - Mojavean desert scrub, Sonoran Desert scrub, Imperial, San Bernardino, San Diego, Riverside; Hayfields Lake	Federal: none  CRPR: 2B.2	5-1310	Aug-Nov	Low – marginal habitat  Not observed
Arizona spurge <i>Chamaesyce arizonica</i>	Perennial herb; Sonoran Desert scrub (sandy); Imperial, Riverside, San Diego; Santa Rosa Mountains	Federal: none  CRPR: 2B.3	50-300	Mar-Apr	Low - distant from known records  Not observed
Flat-seeded spurge <i>Chamaesyce platysperma</i>	Annual herb; Desert dunes - Sonoran Desert scrub (sandy); Imperial Riverside, San Bernardino, San Diego; Coachella Valley	Federal: none  CRPR: 1B.2  BLM sensitive	65-100	Feb-Sep	Low - distant from known records  Not observed
Las Animas colubrina <i>Colubrina californica</i>	Perennial deciduous shrub; Mojavean desert scrub, Sonoran desert scrub Imperial; Riverside, San Diego; Chuckwalla Mountains	Federal: none  CRPR: 2B.3	10-1000	Apr-Jun	Low – marginal habitat  Not observed
Spiny abrojo <i>Condalia globosa</i> var. <i>pubescens</i>	Perennial deciduous shrub, Sonoran desert scrub, Imperial, Riverside, San Diego	Federal: none  CRPR: 4.2	85-1000	Mar-Nov	Low – marginal habitat  Not observed
Foxtail cactus <i>Coryphantha alversonii</i>	perennial stem succulent; sandy or rocky, usually granitic - Mojavean desert scrub, Sonoran desert scrub; Imperial, Riverside, Imperial	Federal: none  CRPR: 4.3	75-1525	Apr-Jun	Low- marginal habitat  Not observed

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Ribbed cryptantha <i>Cryptantha costata</i>	annual herb; sandy - Desert dunes, Mojavean desert scrub, Sonoran desert scrub; Imperial, Inyo, Riverside, San Bernardino, San Diego	Federal: none  CRPR: 4.3	-560	Feb-May	<b>Present</b> – high concentration near northern boundary
Winged cryptantha <i>Cryptantha holoptera</i>	Annual herb; Mojavean desert scrub - Sonoran desert scrub Imperial, Inyo, Riverside, San Bernardino, San Diego; McCoy Mountains	Federal: none  CRPR: 4.3	100-1690	Mar-Apr	Low - distant from known records  Not observed
Wiggins' cholla <i>Cylindropuntia wigginsii</i>	Perennial stem succulent. Sonoran desert scrub (sandy) Imperial, Riverside, San Bernardino, San Diego; Palo Verde	Federal: none  CRPR: 3.3	30-885	Mar	Low - distant from known records  Not observed
Utah milkvine <i>(Funastrum [Cynanchum] utahense)</i>	Perennial herb; sandy or gravelly - Mojavean desert scrub, Sonoran desert scrub; Imperial, Riverside, San Bernardino, San Diego	Federal: none  CRPR: 4.2	100-1435	Mar-Oct	Moderate  Not observed
Glandular ditaxis <i>Ditaxis claryana</i>	perennial herb; sandy; Mojavean desert scrub; Sonoran desert scrub; Imperial, Riverside, San Diego	Federal: none  CRPR: 2B.2	0-465	Oct-Mar	Moderate  Not observed
California ditaxis <i>Ditaxis serrata</i> var. <i>californica</i>	Perennial herb; Sonoran desert scrub; Imperial, Riverside, San Bernardino, San Diego	Federal: none  CRPR: 3.2	30-1000	Mar-Dec	Moderate  Not observed
Cottontop cactus <i>Echinocactus polycephalus</i> var. <i>polycephalus</i>	Perennial stem succulent. Rocky hills, silt valleys; Sonoran desert scrub; Imperial, Inyo, Riverside, San Bernardino, San Diego	Federal: none  CRPR: CBR	<1400	Mar-Aug	Low – marginal habitat  not observed



PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Harwood's Eriastrum <i>Eriastrum harwoodii</i>	annual herb; Desert dunes; Riverside, San Bernardino, San Diego	Federal: none CRPR: 1B.2	125-915	Mar-Jun	<b>Present</b>
California satintail <i>Imperata brevifolia</i>	perennial rhizomatous herb; Chaparral, Coastal scrub, Mojavean desert scrub, Meadows and seeps (often alkali), Riparian scrub; Butte, Fresno, Imperial, Inyo, Kern, Lake, Los Angeles, Orange, Riverside, San Bernardino, Tehama, Tulare, Ventura	Federal: none CRPR: 2B.1	0-1215	Sep-May	Low - distant from known records  Not observed
Pink velvet mallow <i>Horsfordia alata</i>	Perennial shrub; Sonoran desert scrub (rocky); Imperial, Riverside; Palm Springs	Federal: none CRPR: 4.3	100-500	Feb-Dec	Low - distant from known records not observed
Bitter hymenoxys <i>Hymenoxys odorata</i>	Annual herb sandy; Riparian scrub, Sonoran desert scrub; San Bernardino, Riverside, Imperial; near Blythe	Federal: none CRPR: 2B.1	45-150	Feb-Nov	Low - distant from known records Not observed
Spearleaf <i>Matelea parvifolia</i>	Perennial herb; rocky - Mojavean desert scrub, Sonoran desert scrub; Imperial, Riverside, San Bernardino, San Diego; Hayfield Lake	Federal: none CRPR: 2B.3	440-1095	Mar-May	Minimal – no suitable habitat  Not observed
Argus blazing star <i>Mentzelia puberula</i>	Perennial herb; sandy or rocky - Mojavean desert scrub Sonoran desert scrub, Imperial, Riverside, San Bernardino	Federal: none CRPR: 2B.2	90-1280	Mar-May	Moderate  Not observed
Slender wooly heads <i>Nemacaulis denudata var. gracilis</i>	Annual herb; coastal dunes, desert dunes, Sonoran desert scrub; Imperial, Riverside, San Bernardino, San Diego; Arica Mountains	Federal: none CRPR: 2B.2	-450	Mar-May	Low - distant from known records Not observed

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Narrow-leaved sandpaper plant <i>Petalonyx linearis</i>	Perennial shrub; sandy or rocky canyons, generally in creosote bush scrub; Riverside County, Joshua Tree SE Coxcumb Mountains	Federal: none  CRPR: 2B.3	<1000	Mar-May	Minimal – lacks suitable habitat  Not observed
Lobed cherry <i>Physalis lobata</i>	Perennial herb; Mojavean desert scrub (decomposed granitic), Playas; San Bernardino; Hwy 62	Federal: none  CRPR: 2B.3	500-800	May-Jan	Low - distant from known records  Not observed
Desert portulaca <i>Portulaca halimoides</i>	Annual herb; Joshua tree woodland (sandy, San Bernardino, Riverside)	Federal: none  CRPR: 4.2	1000-2000	Sep	Minimal- unsuitable elevation  Not observed
Desert unicorn plant <i>Proboscidea althaeifolia</i>	Perennial herb; gently sloping sandy flats and washes, sometimes roadsides, Sonoran desert scrub; Imperial, Riverside, San Bernardino, San Diego	Federal: none  CRPR: 4.3	85-1000	May-Oct	Moderate  Not observed
Orocopia sage <i>Salvia greatae</i>	Perennial evergreen shrub; Mojavean desert scrub, Sonoran desert scrub; Imperial, Riverside, San Bernardino; Orocopia and Chocolate Mountains	Federal: none CRPR: 1B.3 BLM sensitive	-865	Mar-Apr	Minimal – unsuitable habitat  Not observed
Desert spikemoss <i>Selaginella eremophila</i>	Perennial rhizomatous herb; chaparral, Sonoran desert scrub (gravelly or rocky); Imperial, Riverside, San Diego; Orocopia Mountains	Federal: none  CRPR: 2B.2	200-1295	May-Jul	Minimal - unsuitable  Not observed
Cove's cassia <i>Senna covesii</i>	Perennial herb; dry, sandy desert washes and slopes, Sonoran desert scrub; Imperial, Riverside, Kern, San Bernardino, San Diego	Federal: none  CRPR: 2B.2	225-1295	Mar-Aug	Minimal – unsuitable elevation  Not observed

PLANT SPECIES	FORM; HABITAT; DISTRIBUTION (COUNTIES)	CONSERVATION STATUS	ELEVATION (meters)	BLOOMING PERIOD	POTENTIAL TO OCCUR ON THE PROJECT SITE
Mesquite nest straw <i>Stylocline sonorensis</i>	Annual herb; Sonoran desert scrub (sandy) Known in CA from only a single collection (1930) at Hayfields Dry Lake Possibly extirpated after 1930 by development	Federal: none  CRPR: 2A	+/- 400	Apr	Low - distant from known records  Not observed
Dwarf germander <i>Teucrium cubense ssp. depressum</i>	Annual herb; desert dunes, playas margins; Sonoran desert scrub, Imperial, Riverside; Hayfield Lake	Federal: none  CRPR: 2B.2	45-400	Mar-Nov	Low - distant from known records  Not observed
Jackass clover <i>Wislizenia refracta ssp. refracta</i>	Annual herb; desert dunes, Mojavean desert scrub, playas, sonoran desert scrub, Riverside, San Bernardino	Federal: none  CRPR: 2B.2	600-800	Apr-Nov	Moderate  Not observed
Palmer's jackass clover <i>Wislizenia refracta ssp. Palmeri</i>	perennial deciduous shrub; Chenopod scrub, Desert dunes, Sonoran desert scrub, Sonoran thorn woodland, Riverside, San Diego; Palen sand dunes, Palen Mountains	Federal: none  CRPR: 2B.2	0-300	Jan-Dec	Moderate  Not observed
"Palen Lake atriplex" <i>Atriplex sp. nov. J. Andre</i> ( <i>Atriplex canescens</i> var. <i>macilenta</i> )	Perennial shrub; Saline habitats, playa margins of Palen Dry Lake; Riverside	Federal: none  CRPR: none  BLM sensitive	<160	May-Jun	Low – marginal habitat  Not observed

Federal FE = Federally listed endangered: species in danger of extinction throughout a significant portion of its range

FT = Federally listed, threatened: species likely to become endangered within the foreseeable future

California Rare Plant Rank (CRPR)

CRPR 1A = Presumed extirpated in California and either rare or extinct elsewhere

CRPR 1B = Rare, threatened, or endangered in California and elsewhere

CRPR 2A = Presumed extirpated in California but more common elsewhere

CRPR 2B = Rare, threatened, or endangered in California but more common elsewhere

CRPR 3 = Plants which need more information

CRPR 4 = Limited distribution – a watch list

CBR = Considered, But Rejected

.1 = Seriously endangered in California (high degree/immediacy of threat; over 80% of occurrences threatened)

.2 = Fairly endangered in California (moderate degree/immediacy of threat; 20%-80% of occurrences threatened)

.3 = Not very endangered in California (low degree/immediacy of threats or no current threats known; <20% of occurrences threatened or no current threats known)

#### Bureau of Land Management

BLM Sensitive = may be designated by the BLM California State Director for the following groups of species: proposed or candidate species for listing under the federal ESA, ESA delisted species in the five years following delisting, species listed under the California ESA, California State Species of Special Concern (SSC), California State Fully Protected Species (FP), and California rare plants ranked as List 1B (plants rare, threatened, or endangered in California and elsewhere) identified in the California Department of Fish and Wildlife Special Vascular Plants, Bryophytes, and Lichens List (current version online at <https://www.wildlife.ca.gov/Conservation/Plants/Info>), unless the State Director decides on a case-by-case basis that a particular List 1B species does not warrant sensitive status. (BLM Manual 6840, 2008 and BLM Manual 6840-1, 1996)

## APPENDIX D

### Arica Solar Project Wildlife Species Observed Fall 2019-Spring 2020

COMMON NAME	SCIENTIFIC NAME
<b>Mammals</b>	
Black tailed jackrabbit	<i>Lepus californicus</i>
<b>Burro deer</b>	<b><i>Odocoileus hemionus</i></b>
Coyote	<i>Canis latrans</i>
Desert kangaroo rat	<i>Dipodomys deserti</i>
<b>Desert kit fox</b>	<b><i>Vulpes macrotis</i></b>
Round tail ground squirrel	<i>Xerospermophilus tereticaudus</i>
<b>Reptiles</b>	
Desert horned lizard	<i>Phrynosoma platyrhinos</i>
Desert iguana	<i>Dipsosaurus dorsalis</i>
<b>Desert tortoise</b>	<b><i>Gopherus agassizii</i></b>
Leopard lizard	<i>Gambelia wislizenii</i>
Mohave shovel nosed snake	<i>Chionactis occipitalis</i>
<b>Mojave fringe-toed lizard</b>	<b><i>Uma scoparia</i></b>
Side blotched lizard	<i>Uta stansburiana</i>
Western whiptail lizard	<i>Aspidoscelis tigris</i>
Zebra-tailed lizard	<i>Calisaurus draconoides</i>
<b>Birds</b>	
American kestrel	<i>Falco sparverius</i>
American pipit	<i>Anthus rubescens</i>
Anna's hummingbird	<i>Calypte anna</i>
Ash-throated flycatcher	<i>Myiarchus cinerascens</i>
Barn Swallow	<i>Hirundo rustica</i>
Black phoebe	<i>Sayornis nigricans</i>
Black tailed gnatcatcher	<i>Poliophtila melanura</i>
Black throated sparrow	<i>Amphispiza bilineata</i>
Blue grey gnatcatcher	<i>Poliophtila caerulea</i>
Brewer's sparrow	<i>Spizella breweri</i>
<b>Burrowing owl</b>	<b><i>Athene cunicularia</i></b>

COMMON NAME	SCIENTIFIC NAME
Cactus wren	<i>Campylorhynchus brunneicapillus</i>
Cliff swallow	<i>Petrochelidon pyrrhonota</i>
Common poorwill	<i>Phalaenoptilus nuttallii</i>
Common raven	<i>Corvus corax</i>
Cooper's hawk	<i>Accipiter cooperii</i>
European starling	<i>Sturnus vulgaris</i>
Gambel's quail	<i>Callipepla gambelii</i>
Greater roadrunner	<i>Geococcyx californianus</i>
Great-tailed grackle	<i>Quiscalus mexicanus</i>
Horned lark	<i>Eremophila alpestris</i>
House finch	<i>Carpodacus menicanus</i>
House wren	<i>Troglodytes aedon</i>
<b>LeConte's Thrasher</b>	<b><i>Toxostoma lecontei</i></b>
Lesser goldfinch	<i>Spinus psaltria</i>
Lesser nighthawk	<i>Chordeiles acutipennis</i>
<b>Loggerhead shrike</b>	<b><i>Lanius ludovicianus</i></b>
Mallard	<i>Anas platyrhynchos</i>
Mourning dove	<i>Zenaida macroura</i>
Northern flicker	<i>Colaptes auratus</i>
Northern harrier	<i>Circus cyaneus</i>
Northern mockingbird	<i>Mimus polyglottos</i>
Northern rough-winged swallow	<i>Stelgidopteryx serripennis</i>
Osprey	<i>Pandio haliaetus</i>
<b>Prairie falcon</b>	<b><i>Falco mexicanus</i></b>
Red tailed hawk	<i>Buteo jamaicensis</i>
Rock wren	<i>Salpinctes obsoletus</i>
Ruby crowned kinglet	<i>Regulus calendula</i>
Sage thrasher	<i>Oreoscoptes montanus</i>
Sagebrush sparrow	<i>Artemisiospiza nevadensis</i>
Say's phoebe	<i>Sayornis saya</i>
Short-eared owl	<i>Asio flammeus</i>
Snow goose	<i>Chen caerulescens</i>
<b>Swainson's hawk</b>	<b><i>Buteo swainsoni</i></b>
Tree swallow	<i>Tachycineta bicolor</i>
Turkey vulture	<i>Cathartes aura</i>
Verdin	<i>Auriparus flaviceps</i>
Violet green swallow	<i>Tachycineta thalassina</i>
Western meadowlark	<i>Sturnella neglecta</i>



COMMON NAME	SCIENTIFIC NAME
White-crowned sparrow	<i>Zonotrichia leucophrys</i>
Yellow-rumped (Audubon's)warbler	<i>Setophaga coronata</i>

**BOLD** = special status species

## APPENDIX E

### Arica Solar Project Plants Species Observed Fall 2019-Spring 2020

FAMILY	SCIENTIFIC NAME	COMMON NAME
Amaranthaceae	<i>Tidestromia suffruticosa</i> var. <i>oblongifolia</i>	Honeysweet
Asteraceae	<i>Ambrosia dumosa</i>	White bursage
Asteraceae	<i>Ambrosia salsola</i>	Cheesebush
Asteraceae	<i>Atrichocercis platyphylla</i>	Gravel ghost
Asteraceae	<i>Baileya pauciradiata</i>	Lax flower
Asteraceae	<i>Baileya pleniradiata</i>	Wooly marigold
Asteraceae	<i>Bebbia juncea</i> var. <i>aspera</i>	Rush sweetbush
Asteraceae	<i>Calycoseris wrightii</i>	White tackstem
Asteraceae	<i>Chaenactis carphoclinia</i>	Pebble pincushion
Asteraceae	<i>Chaenactis fremontii</i>	Fremont's pincushion
Asteraceae	<i>Chaenactis stevioides</i>	Desert pincushion
Asteraceae	<i>Encelia farinosa</i>	Brittlebush
Asteraceae	<i>Encelia frutescens</i>	Button brittlebush
Asteraceae	<i>Geraea canescens</i>	Desert sunflower
Asteraceae	<i>Logfia depressa</i>	Dwarf cottonrose
Asteraceae	<i>Malacothrix glabrata</i>	Desert dandelion
Asteraceae	<i>Monoptilon bellioides</i>	Mojave desert star
Asteraceae	<i>Palafoxia arida</i> var. <i>arida</i>	Spanish needle
Asteraceae	<i>Perityle emoryi</i>	Emory's rockdaisy
Asteraceae	<i>Psathyrotes ramosissima</i>	Turtleback
Asteraceae	* <i>Sonchus oleraceus</i>	Sow thistle
Asteraceae	<i>Stephanomeria pauciflora</i>	Wire lettuce
Boraginaceae	<i>Amsinckia tessellata</i>	Devil's lettuce
Boraginaceae	<i>Cryptantha angustifolia</i>	Narrow leaved cryptantha
<b>Boraginaceae</b>	<b><i>Cryptantha costata</i></b>	<b>Ashen forget-me-not</b>
Boraginaceae	<i>Cryptantha maritima</i>	Guadalupe cryptantha
Boraginaceae	<i>Cryptantha nevadensis</i>	Nevada forget me not
Boraginaceae	<i>Cryptantha pterocarya</i>	Winged nut forget-me-not
Boraginaceae	<i>Cryptantha utahensis</i>	Scented forget-me-not
Boraginaceae	<i>Eucrypta micrantha</i>	Desert eucrypta
Boraginaceae	<i>Nama demissa</i> var. <i>demissa</i>	Purple mat

<b>FAMILY</b>	<b>SCIENTIFIC NAME</b>	<b>COMMON NAME</b>
Boraginaceae	<i>Nama hispida</i> var. <i>spathulata</i>	Purplemat
Boraginaceae	<i>Pectocarya heterocarpa</i>	Cuckwalla pectocarya
Boraginaceae	<i>Pectocarya platycarpa</i>	Broad nutted comb-bur
Boraginaceae	<i>Phacelia crenulata</i>	Notch leaved phacelia
Boraginaceae	<i>Phacelia crenulata</i> var. <i>ambigua</i>	Purplestem phacelia
Boraginaceae	<i>Phacelia crenulata</i> var. <i>crenulata</i>	Heliotrope phacelia
Boraginaceae	<i>Phacelia distans</i>	Common phacelia
Boraginaceae	<i>Phacelia tanacetifolia</i>	Tansy leafed phacelia
Boraginaceae	<i>Tiquilia plicata</i>	Fanleaf crinklemat
Brassicaceae	* <i>Brassica tournefortii</i>	Sahara mustard
Brassicaceae	<i>Caulanthus lasiophyllus</i>	California mustard
Brassicaceae	<i>Descurainia pinnata</i>	Western tansy mustard
Brassicaceae	<i>Dithyrea californica</i>	Spectacle pod
Brassicaceae	<i>Lepidium lasiocarpum</i>	pepperweed
Brassicaceae	* <i>Sisymbrium irio</i>	London rocket
Brassicaceae	* <i>Sisymbrium orientale</i>	Hedge mustard
Cactaceae	<i>Cylindropuntia echinocarpa</i>	Golden cholla
Cactaceae	<i>Cylindropuntia ramosissima</i>	Diamond cholla
Caryophyllaceae	<i>Achyronychia cooperi</i>	Frost mat
Chenopodiaceae	<i>Atriplex canescens</i>	Hoary saltbush
Chenopodiaceae	<i>Atriplex polycarpa</i>	Allscale saltbush
Chenopodiaceae	* <i>Salsola tragus</i>	Russian thistle
Convolvulaceae	<i>Cucscuta</i> sp.	Dodder
Convolvulaceae	<i>Phoradendron californicum</i>	California mesquite mistletoe
Cucurbitaceae	<i>Brandegea bigelovii</i>	Desert starvine
Euphorbiaceae	<i>Croton californicus</i>	California croton
Euphorbiaceae	<i>Ditaxis lanceolata</i>	Narrowleaf ditaxis
Euphorbiaceae	<i>Ditaxis neomexicana</i>	New Mexico ditaxis
Euphorbiaceae	<i>Euphorbia polycarpa</i>	Smallseed sandmat
Euphorbiaceae	<i>Stillingia spinulosa</i>	Broad leaved stillingia
Fabaceae	<i>Acmispon strigosus</i>	Strigose lotus
Fabaceae	<i>Astragalus aridus</i>	Annual desert milk vetch
Fabaceae	<i>Astragalus didymocarpus</i>	Dwarf white milk vetch
<b>Fabaceae</b>	<b><i>Astragalus insularis</i> var. <i>harwoodii</i></b>	<b>Harwood's milk-vetch</b>

<b>FAMILY</b>	<b>SCIENTIFIC NAME</b>	<b>COMMON NAME</b>
Fabaceae	<i>Dalea mollissima</i>	Silky dalea
Fabaceae	<i>Lupinus arizonicus</i>	Arizona lupine
Fabaceae	<i>Lupinus shockleyi</i>	Shockley lupine
Fabaceae	<i>Marina parryi</i>	Parry's false prairie clover
Fabaceae	<i>Olneya tesota</i>	Desert ironwood
Fabaceae	<i>Parkinsonia florida</i>	Blue palo verde
Fabaceae	<i>Prosopis glandulosa</i> var. <i>torreyana</i>	Honey mesquite
Fabaceae	<i>Psoralea arguta</i>	Indigo bush
Fabaceae	<i>Psoralea schottii</i>	Schott's indigo bush
Fabaceae	<i>Psoralea spinosa</i>	Smoke tree
Fabaceae	<i>Senegalia greggii</i>	Catclaw acacia
Geraniaceae	* <i>Erodium cicutarium</i>	Coastal heron's bill
Geraniaceae	<i>Erodium texanum</i>	Desert heron's bill
Lamiaceae	<i>Condea emoryi</i> (= <i>Hyptis emoryi</i> )	Desert lavender
Lamiaceae	<i>Salvia columbariae</i>	Chia sage
Liliaceae	<i>Hesperocallis undulata</i>	Desert lily
Loasaceae	<i>Mentzelia albicaulis</i>	White stemmed stickleaf
Loasaceae	<i>Mentzelia involucrata</i>	Whitebract blazingstar
Malvaceae	<i>Eremalche exilis</i>	White mallow
Malvaceae	<i>Eremalche rotundifolia</i>	Desert fivespot
Malvaceae	<i>Hibiscus denudatus</i>	Paleface
Malvaceae	<i>Sphaeralcea ambigua</i>	Desert globemallow
Nyctaginaceae	<i>Abronia villosa</i> var. <i>villosa</i>	Hairy sand verbena
Nyctaginaceae	<i>Boerhavia wrightii</i>	Wright's boerhavia
Nyctaginaceae	<i>Mirabilis laevis</i>	Desert wishbone bush
Onagraceae	<i>Chylismia brevipes</i> subsp. <i>brevipes</i>	Golden suncup
Onagraceae	<i>Chylismia claviformis</i>	Browneyes
Onagraceae	<i>Eremothera boothii</i> subsp. <i>condensata</i>	Booth's suncup
Onagraceae	<i>Eremothera refracta</i>	Narrow leaved primrose
Onagraceae	<i>Oenothera caespitosa</i>	Fragrant evening primrose
Onagraceae	<i>Oenothera deltoides</i> subsp. <i>deltoides</i>	Birdcage desert primrose
Onagraceae	<i>Oenothera primiveris</i>	Yellow desert evening primrose
Papaveraceae	<i>Argemone munita</i>	Prickly poppy

<b>FAMILY</b>	<b>SCIENTIFIC NAME</b>	<b>COMMON NAME</b>
Papaveraceae	<i>Eschscholzia glyptosperma</i>	Desert gold poppy
Papaveraceae	<i>Eschscholzia minutiflora</i>	Small flowered poppy
Papaveraceae	<i>Eschscholzia parishii</i>	Pygmy poppy
Plantaginaceae	<i>Plantago ovata</i>	Wooly plantain
Poaceae	<i>Aristida adscensionis</i>	Three-awn
Poaceae	<i>Aristida californica subsp. californica</i>	California three-awn
Poaceae	<i>Aristida purpurea</i>	Purple three-awn
Poaceae	* <i>Bromus tectorum</i>	Downy chess
Poaceae	<i>Festuca octoflora</i>	Fescue
Poaceae	<i>Hilaria rigida</i>	Big galleta grass
Poaceae	* <i>Schismus barbatus</i>	Common mediterranean grass
<b>Polemoniaceae</b>	<b><i>Eriastrum harwoodii</i></b>	<b>Harwood's eriastrum</b>
Polemoniaceae	<i>Gilia stellata</i>	Star gilia
Polemoniaceae	<i>Linanthus jonesii</i>	Jones linanthus
Polemoniaceae	<i>Loeseliastrum matthewsii</i>	Desert calico
Polemoniaceae	<i>Loeseliastrum schottii</i>	Schott gilia
Polygonaceae	<i>Chorizanthe brevicornu</i>	Brittle spineflower
Polygonaceae	<i>Chorizanthe rigida</i>	Devil's spineflower
Polygonaceae	<i>Eriogonum inflatum</i>	Desert trumpet
Polygonaceae	<i>Eriogonum thomasii</i>	Thomas' buckwheat
Resedaceae	<i>Oligomeris linifolia</i>	Leaved cambess
<b>Simaroubaceae</b>	<b><i>Castela emoryi</i></b>	<b>Crucifixion thorn</b>
Solanaceae	<i>Datura discolor</i>	Small datura
Solanaceae	<i>Datura wrightii</i>	Jimson weed
Solanaceae	<i>Lycium andersonii</i>	Anderson's desert thorn
Solanaceae	<i>Physalis crassifolia</i>	Ground cherry
Zygophyllaceae	<i>Fagonia laevis</i>	California fagonia
Zygophyllaceae	<i>Larrea tridentata</i>	Creosote bush

**BOLD** = special status species

\*= non-native species