# Appendix F

Soils Report



United States Department of Agriculture

Natural Resources


Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

# Custom Soil Resource Report for Tulare County, Western Part, California

**Carleton Acres Development** 



# Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2\_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

# Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	
Soil Map	
Legend	
Map Unit Legend	
Map Unit Descriptions	
Tulare County, Western Part, California	
101—Akers-Akers, saline-Sodic, complex, 0 to 2 percent slopes	13
122—Grangeville sandy loam, drained, 0 to 2 percent slopes	15
Soil Information for All Uses	18
Suitabilities and Limitations for Use	18
Land Classifications	18
California Revised Storie Index (CA) (Carleton Acres)	18
Irrigated Capability Class (Carleton Acres)	
References	27

# **How Soil Surveys Are Made**

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

# Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



	MAP LEGEND			MAP INFORMATION	
Area of Int	Area of Interest (AOI) Area of Interest (AOI)		Spoil Area	The soil surveys that comprise your AOI were mapped at	
			Stony Spot	1:24,000.	
Soils	Soil Map Unit Polygons	Ø	Very Stony Spot	Warning: Soil Map may not be valid at this scale.	
	Soil Map Unit Lines	Ŷ	Wet Spot		
~	Soil Map Unit Points	$\triangle$	Other	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil	
			Special Line Features	line placement. The maps do not show the small areas of	
Special (0)	Point Features Blowout	Water Fea	atures	contrasting soils that could have been shown at a more detailed scale.	
×	Borrow Pit	$\sim$	Streams and Canals		
×	Clay Spot	Transport		Please rely on the bar scale on each map sheet for map	
ô	Closed Depression	+++	Rails	measurements.	
×	Gravel Pit	~	Interstate Highways	Source of Map: Natural Resources Conservation Service	
°.	Gravelly Spot	Coordinate SV	Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)		
	Landfill	~	Major Roads		
Ň.	Lava Flow	~	Local Roads	Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts	
<u>بلد</u>	Marsh or swamp	Backgrou	ground Aerial Photography	distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more	
~	Mine or Quarry	No.		accurate calculations of distance or area are required.	
Ô	Miscellaneous Water			This product is generated from the LISDA NDCS partified data as	
Ő	Perennial Water			This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.	
Š	Rock Outcrop			Quill Querran Anna Talana Querra Manter Data Quillantia	
* +	Saline Spot			Soil Survey Area: Tulare County, Western Part, California Survey Area Data: Version 15, Sep 3, 2021	
	Sandy Spot				
	Severely Eroded Spot			Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.	
0	Sinkhole			Data(a) agrial imagaa wara nhatagranhadi Jan 20, 2024	
>	Slide or Slip			Date(s) aerial images were photographed: Jan 30, 2021—Feb 6, 2021	
ø	Sodic Spot			The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor	
				compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.	

## **Map Unit Legend**

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
101	Akers-Akers, saline-Sodic, complex, 0 to 2 percent slopes	268.5	53.3%
122	Grangeville sandy loam, drained, 0 to 2 percent slopes	235.1	46.7%
Totals for Area of Interest	·	503.6	100.0%

# **Map Unit Descriptions**

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the

development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

### Tulare County, Western Part, California

#### 101—Akers-Akers, saline-Sodic, complex, 0 to 2 percent slopes

#### **Map Unit Setting**

National map unit symbol: hp6z
Elevation: 230 to 350 feet
Mean annual precipitation: 8 to 12 inches
Mean annual air temperature: 63 to 64 degrees F
Frost-free period: 225 to 300 days
Farmland classification: Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season

#### **Map Unit Composition**

Akers and similar soils: 60 percent Akers, saline-sodic, and similar soils: 25 percent Minor components: 15 percent Estimates are based on observations, descriptions, and transects of the mapunit.

#### **Description of Akers**

#### Setting

Landform: Fan remnants Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Side slope Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium derived from granitic rock sources

#### **Typical profile**

*Ap - 0 to 16 inches:* fine sandy loam *Bk - 16 to 60 inches:* fine sandy loam

#### **Properties and qualities**

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Negligible
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Very rare
Frequency of ponding: None
Calcium carbonate, maximum content: 10 percent
Gypsum, maximum content: 2 percent
Maximum salinity: Nonsaline to slightly saline (0.0 to 4.0 mmhos/cm)
Sodium adsorption ratio, maximum: 12.0
Available water supply, 0 to 60 inches: High (about 9.7 inches)

#### Interpretive groups

Land capability classification (irrigated): 1 Land capability classification (nonirrigated): 4c Hydrologic Soil Group: B Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### **Description of Akers, Saline-sodic**

#### Setting

Landform: Fan remnants Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Side slope Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium derived from granitic rock sources

#### **Typical profile**

*Ap - 0 to 15 inches:* fine sandy loam *Bk - 15 to 60 inches:* fine sandy loam

#### **Properties and qualities**

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Negligible
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Very rare
Frequency of ponding: None
Calcium carbonate, maximum content: 10 percent
Gypsum, maximum content: 2 percent
Maximum salinity: Slightly saline to moderately saline (4.0 to 8.0 mmhos/cm)
Sodium adsorption ratio, maximum: 30.0
Available water supply, 0 to 60 inches: High (about 9.1 inches)

#### Interpretive groups

Land capability classification (irrigated): 2s Land capability classification (nonirrigated): 4s Hydrologic Soil Group: C Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### **Minor Components**

#### Colpien

Percent of map unit: 3 percent Landform: Fan remnants Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Tujunga

Percent of map unit: 3 percent Landform: Flood plains Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Tagus

Percent of map unit: 2 percent Landform: Fan remnants Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Hanford

Percent of map unit: 2 percent Landform: Alluvial fans, flood plains Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Yettem

Percent of map unit: 2 percent Landform: Alluvial fans, flood plains Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Grangeville

Percent of map unit: 2 percent Landform: Alluvial fans, flood plains Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: No

#### Unnamed, ponded

Percent of map unit: 1 percent Landform: Depressions Ecological site: R017XY907CA - Aridic Alkali Desert Hydric soil rating: Yes

#### 122—Grangeville sandy loam, drained, 0 to 2 percent slopes

#### Map Unit Setting

National map unit symbol: hp4s Elevation: 190 to 400 feet Mean annual precipitation: 8 to 12 inches Mean annual air temperature: 63 to 64 degrees F Frost-free period: 250 to 275 days Farmland classification: Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season

#### Map Unit Composition

Grangeville and similar soils: 90 percent Minor components: 10 percent Estimates are based on observations, descriptions, and transects of the mapunit.

#### **Description of Grangeville**

#### Setting

Landform: Alluvial fans, flood plains Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Base slope Down-slope shape: Linear Across-slope shape: Convex, linear Parent material: Alluvium derived from granitic rock sources

#### **Typical profile**

*Ap - 0 to 16 inches:* sandy loam *Bg - 16 to 27 inches:* sandy loam *2C - 27 to 67 inches:* stratified loamy sand to silt loam

#### **Properties and qualities**

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Negligible
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Rare
Frequency of ponding: None
Calcium carbonate, maximum content: 5 percent
Gypsum, maximum content: 1 percent
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water supply, 0 to 60 inches: Moderate (about 6.6 inches)

#### Interpretive groups

Land capability classification (irrigated): 1 Land capability classification (nonirrigated): 4c Hydrologic Soil Group: A Ecological site: R017XY906CA - Non-Alkali San Joaquin Valley Desert Hydric soil rating: Yes

#### **Minor Components**

#### Yettem

Percent of map unit: 3 percent Landform: Flood plains, alluvial fans Ecological site: R017XY904CA - Subirrigated Deep Alluvial Fans Hydric soil rating: No

#### Tujunga

Percent of map unit: 3 percent Landform: Flood plains Ecological site: R017XY904CA - Subirrigated Deep Alluvial Fans Hydric soil rating: No

#### Grangeville, saline-sodic

Percent of map unit: 2 percent Landform: Alluvial fans, flood plains Ecological site: R017XY904CA - Subirrigated Deep Alluvial Fans Hydric soil rating: Yes

#### Nord

Percent of map unit: 1 percent Landform: Alluvial fans, flood plains Ecological site: R017XY904CA - Subirrigated Deep Alluvial Fans Hydric soil rating: No

#### Hanford

*Percent of map unit:* 1 percent *Landform:* Alluvial fans, flood plains

*Ecological site:* R017XY904CA - Subirrigated Deep Alluvial Fans *Hydric soil rating:* No

# **Soil Information for All Uses**

### Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

### Land Classifications

Land Classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

### California Revised Storie Index (CA) (Carleton Acres)

The Revised Storie Index is a rating system based on soil properties that govern the potential for soil map unit components to be used for irrigated agriculture in California.

The Revised Storie Index assesses the productivity of a soil from the following four characteristics:

- Factor A: degree of soil profile development
- Factor B: texture of the surface layer
- Factor C: steepness of slope

- Factor X: drainage class, landform, erosion class, flooding and ponding frequency and duration, soil pH, soluble salt content as measured by electrical conductivity, and sodium adsorption ratio

Revised Storie Index numerical ratings have been combined into six classes as follows:

- Grade 1: Excellent (81 to 100)
- Grade 2: Good (61 to 80)
- Grade 3: Fair (41 to 60)
- Grade 4: Poor (21 to 40)
- Grade 5: Very poor (11 to 20)
- Grade 6: Nonagricultural (10 or less)

The components listed for each map unit in the accompanying Summary by Map Unit table in Web Soil Survey or the Aggregation Report in Soil Data Viewer are determined by the aggregation method chosen. An aggregated rating class is shown for each map unit. The components listed for each map unit are only those that have the same rating class as the one shown for the map unit. The percent composition of each component in a particular map unit is given to help the user better understand the extent to which the rating applies to the map unit.

Other components with different ratings may occur in each map unit. The ratings for all components, regardless the aggregated rating of the map unit, can be viewed by generating the equivalent report from the Soil Reports tab in Web Soil Survey or from the Soil Data Mart site. Onsite investigation may be needed to validate these interpretations and to confirm the identity of the soil on a given site.



	MAP LEGEND			MAP INFORMATION	
Area of Interest (AOI) Area of Interest (AOI)		Grade 5 - Very Poor		The soil surveys that comprise your AOI were mapped at 1:24,000.	
Soils			Grade 6 - Nonagricultural		
	ting Polygons		Not rated	Warning: Soil Map may not be valid at this scale.	
	Grade 1 - Excellent		Not rated or not available	Entergoment of more beyond the cools of morning concerned	
	Grade 2 - Good	Water Fea		Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil	
	Grade 3 - Fair	$\sim$	Streams and Canals	line placement. The maps do not show the small areas of	
	Grade 4 - Poor	Transportation		contrasting soils that could have been shown at a more detailed scale.	
	Grade 5 - Very Poor	+++	Rails		
	,	~	Interstate Highways	Please rely on the bar scale on each map sheet for map	
	Grade 6 - Nonagricultural	~	US Routes	measurements.	
	Not rated	$\sim$	Major Roads	Source of Map: Natural Resources Conservation Service	
	Not rated or not available	~	Local Roads	Web Soil Survey URL:	
Soil Ra	Soil Rating Lines		nd	Coordinate System: Web Mercator (EPSG:3857)	
~	Grade 1 - Excellent	Mar.	Aerial Photography	Maps from the Web Soil Survey are based on the Web Mercato	
~	Grade 2 - Good			projection, which preserves direction and shape but distorts	
~	Grade 3 - Fair			distance and area. A projection that preserves area, such as th	
~	Grade 4 - Poor			Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.	
~	Grade 5 - Very Poor				
	Grade 6 - Nonagricultural			This product is generated from the USDA-NRCS certified data a of the version date(s) listed below.	
	Not rated			of the version date(s) listed below.	
				Soil Survey Area: Tulare County, Western Part, California	
	Not rated or not available			Survey Area Data: Version 15, Sep 3, 2021	
Soil Ra	Crode 1 Excellent			Soil map units are labeled (as space allows) for map scales	
	Grade 1 - Excellent			1:50,000 or larger.	
	Grade 2 - Good			Date(a) parial images were photographed. Let 20, 2024 Fel	
	Grade 3 - Fair			Date(s) aerial images were photographed: Jan 30, 2021—Fel 2021	
	Grade 4 - Poor				
				The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.	

### Table—California Revised Storie Index (CA) (Carleton Acres)

Map unit symbol	Map unit name	Rating	Component name (percent)	Acres in AOI	Percent of AOI
101	Akers-Akers, saline- Sodic, complex, 0 to 2 percent slopes	Grade 1 - Excellent	Akers (60%) Akers, saline-sodic (25%)	268.5	53.3%
122	Grangeville sandy loam, drained, 0 to 2 percent slopes	Grade 2 - Good	Grangeville (90%)	235.1	46.7%
Totals for Area of Interest				503.6	100.0%

# Rating Options—California Revised Storie Index (CA) (Carleton Acres)

Aggregation Method: Dominant Condition Component Percent Cutoff: None Specified Tie-break Rule: Lower

### Irrigated Capability Class (Carleton Acres)

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations that show suitability and limitations of groups of soils for rangeland, for woodland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels-capability class, subclass, and unit. Only class and subclass are included in this data set.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

Class 1 soils have few limitations that restrict their use.

Class 2 soils have moderate limitations that reduce the choice of plants or that require moderate conservation practices.

Class 3 soils have severe limitations that reduce the choice of plants or that require special conservation practices, or both.

Class 4 soils have very severe limitations that reduce the choice of plants or that require very careful management, or both.

Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.


Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.



MAP INFORMATION

#### MAP LEGEND



### Table—Irrigated Capability Class (Carleton Acres)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
101	Akers-Akers, saline- Sodic, complex, 0 to 2 percent slopes	1	268.5	53.3%
122	Grangeville sandy loam, drained, 0 to 2 percent slopes	1	235.1	46.7%
Totals for Area of Intere	st	503.6	100.0%	

### Rating Options—Irrigated Capability Class (Carleton Acres)

Aggregation Method: Dominant Condition Component Percent Cutoff: None Specified Tie-break Rule: Higher

# References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2\_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2\_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2\_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcs142p2\_052290.pdf